

**BOARD OF DENTISTRY
COUNCIL ON DENTAL HYGIENE
AGENDA**

**Call In Number (888) 670-3525, Participant Pass Code: 4552635641
December 7, 2016
6:00 p.m. EST**

Participants in this public meeting should be aware that these proceedings are being recorded and that an audio file of the meeting will be posted to the board's website.

I. CALL TO ORDER

II. DISCUSSION ITEMS

a. Silver Diamine Fluoride

- i. Silver Diamine Fluoride in Caries Management Article**
- ii. Silver Diamine Fluoride 38% Scientific Literature Review**
- iii. Breakthrough Therapy Designation for Silver Diamine Fluoride Granted by the Food and Drug Administration Article**
- iv. Email from Dr. Scott Tomar, Professor & Interim Chair, university of Florida**

b. Dental Hygiene Curriculum Framework

III. OLD BUSINESS

IV. NEW BUSINESS

***To connect to the conference call, dial the following number: 888-670-3525 a minute or two prior to the start time of the meeting. You will then be prompted to enter a "participant pass code", which is 4552635641, followed by the # sign. Please mute your line after joining the call.**



Science in the News

Silver Diamine Fluoride in Caries Management

July 12, 2016

A recent article in the *New York Times* highlighted use of silver diamine fluoride (SDF) as an alternative approach to treatment of cavities in children.¹ The article highlighted that SDF was faster and cheaper than drilling and filling; and it mentioned the downside that when applied, SDF blackens the tooth.

Silver diamine fluoride (SDF) is a colorless liquid that at pH 10 is 24.4% to 28.8% (weight/volume) silver and 5.0% to 5.9% fluoride.² Just as 5% sodium fluoride varnish has FDA clearance as a Class II medical device for the treatment of tooth hypersensitivity, FDA classified SDF as a fluoride and cleared its use as a Class II medical device for the same indication. A number of products are currently available in other countries, but at this time, Advantage Arrest™ (Elevate Oral Care, L.L.C.) is the only commercially available SDF product for dental treatment in the U.S.³

Although FDA-cleared for use in the management of hypersensitivity, SDF received coverage in the *Times* for its use in treating cavities in children, although this might be more accurately described as caries control and management. Likely a result of its fluoride content, when applied to a carious lesion, SDF has also been shown to lower caries risk of the adjacent tooth surface.⁴ While the *Times* article focused on the use of SDF in young children, it has also been shown to be effective in management of root caries in the elderly.^{5,6} It likely has additional applicability as an interim approach for managing problematic caries in individuals currently unable to tolerate more involved dental treatment.

SDF is not a complete solution to caries risk. Single application has been reported to be insufficient for sustained benefit.⁷ Its downsides include a reportedly unpleasant metallic taste, potential to irritate gingival and mucosal surfaces, and the characteristic black staining of the tooth surfaces to which it is applied.³

A search of ClinicalTrials.gov for silver diamine fluoride returned 14 studies; 7 of which have been completed, 5 that are recruiting, and 2 that have not yet begun recruiting.⁸ This suggests that more scientific insight and news will likely be forthcoming about this product.

References

1. Saint Louis C. A Cavity-Fighting Liquid Lets Kids Avoid Dentists' Drills. *New York Times* 2016. <http://www.nytimes.com/2016/07/12/health/silver-diamine-fluoride-dentist-cavities.html>. Accessed July 11, 2016.

2. Mei ML, Chu CH, Lo EC, Samaranayake LP. Fluoride and silver concentrations of silver diammine fluoride solutions for dental use. *Int J Paediatr Dent* 2013;23(4):279-85.
 3. Mei ML, Lo EC, Chu CH. Clinical Use of Silver Diamine Fluoride in Dental Treatment. *Compend Contin Educ Dent* 2016;37(2):93-8; quiz100.
 4. Llodra JC, Rodriguez A, Ferrer B, et al. Efficacy of silver diamine fluoride for caries reduction in primary teeth and first permanent molars of schoolchildren: 36-month clinical trial. *J Dent Res* 2005;84(8):721-4.
 5. Li R, Lo EC, Liu BY, Wong MC, Chu CH. Randomized clinical trial on arresting dental root caries through silver diammine fluoride applications in community-dwelling elders. *J Dent* 2016.
 6. Zhang W, McGrath C, Lo EC, Li JY. Silver diamine fluoride and education to prevent and arrest root caries among community-dwelling elders. *Caries Res* 2013;47(4):284-90.
 7. Horst JA, Ellenikiotis H, Milgrom PL. UCSF Protocol for Caries Arrest Using Silver Diamine Fluoride: Rationale, Indications and Consent. *J Calif Dent Assoc* 2016;44(1):16-28.
 8. U.S. National Institutes of Health. ClinicalTrials.gov search for silver diamine fluoride. <https://clinicaltrials.gov/ct2/results?term=silver+diamine+fluoride&Search=Search>. Accessed July 11, 2016.
-

About Science in the News

Science in the News is a service by the American Dental Association (ADA) to its members to present current information about science topics in the news. The ADA is a professional association of dentists committed to the public's oral health, ethics, science and professional advancement; leading a unified profession through initiatives in advocacy, education, research and the development of standards. As a science-based organization, the ADA's evaluation of the scientific evidence may change as more information becomes available. Your thoughts would be greatly appreciated.

Content on ADA.org is for informational purposes only, is neither intended to and does not establish a standard of care, and is not a substitute for professional judgment, advice, diagnosis, or treatment. ADA is not responsible for information on external websites linked to this website.

Copyright © 2016 American Dental Association. Reproduction or republication strictly prohibited without prior written permission.

Silver Diamine Fluoride 38%

Scientific Literature Review

April 2016

Silver Diamine Fluoride (SDF) 38% has been receiving a great deal of attention by U.S. dental professionals since it was cleared for use by the Food and Drug Administration in August 2104 under the provisions of the Federal Food, Drug and Cosmetics Act. The Cleared Indications For Use are for the “Treatment of dentinal hypersensitivity. For use in adults over the age of 21.”

In the age of the Internet, access to information that can sometimes be credible and sometimes not, could cause confusion about the history, safety and efficacy of SDF. In addition, a number of local television news programs and social media postings around the U.S. have recently begun communicating information about the use of SDF by both general and pediatric dentists who have begun using it for the treatment of carious lesions in populations of all ages.

While SDF only recently received FDA Clearance it has been used by dental professionals outside the U.S. for both the treatment of dentinal hypersensitivity and as a caries therapy for more than 45 years. This review is intended to provide U.S medical professionals with an understanding of the history of SDF around the world, including the most current information available regarding its use in the U.S.

Under federal law, the use of a drug or medical device by a licensed medical professional for an indication not Approved or Cleared by the FDA is allowable and not uncommon. This is termed “off-label” use.

As the organization permitted to market the only FDA Cleared SDF product in the United States, (Advantage Arrest™ Silver Diamine Fluoride 38%), it is our intention to provide a review of all scientific literature available to us in order to help insure that medical professionals, and through them, their patients are as well informed as possible about this therapy.

This document is not assumed to contain all published information regarding SDF, as that would be virtually impossible, since SDF has been in use in many countries around the world for decades. It is however meant to provide a fair and balanced view of the benefits and risks of the use of SDF. If, after reading this document, you have any questions please send an email to the address below and we will get back to you promptly.

Please address any questions to:

Steve Pardue
Elevate Oral Care
spardue@elevateoralcare.com

Table of Contents

Advantage Arrest Package Insert	3
The Short Term Effects of Diammine Silver Fluoride on Tooth Sensitivity: a Randomized Controlled Trial	4
J Dent Res 90(2):203-208, 2011 J.L. Castillo ¹ , S. Rivera, T. Aparicio, R. Lazo ¹ , T.-C. Aw, L.L. Mancl, and P. Milgrom	
UCSF Protocol for Caries Arrest Using Silver Diamine Fluoride: Rationale, Indications and Consent	10
CDA Journal , Vol 44, No.1 Jeremy A. Horst, DDS, PhD; Hellene Ellenikiotis, DDS; and Peter L. Milgrom, DDS	
Clinical Use of Silver Diamine Fluoride in Dental Treatment	22
Compendium of Continuing Education Volume 37, Number 2 May L. Mei, BDS, MDS, PhD; Edward Chin-Man Lo, BDS, MDS, PhD; and Chun-Hung Chu, BDS, MDS, PhD.	
Fresh Approach to Caries Arrest in Adults	23
Decisions in Dentistry Volume 1, Number 1 Dr. John Featherstone, Dean of the University of California San Francisco School of Dentistry and Dr. Jeremy Horst, DDS, PhD	
Additional Articles of Interest	24
Frequently Asked Questions	31

Advantage Arrest Package Insert

Advantage Arrest™

Silver Diamine Fluoride 38%

Professional Tooth Desensitizer

Rx Only

Desensitizing Ingredient: Aqueous Silver Diamine Fluoride, 38.3% to 43.2% w/v

Inactive Ingredients: Purified water

Clinical Pharmacology: Product forms insoluble precipitates with calcium or phosphate in the dentinal tubules to block nerve impulses.

Indication and Usage: Treatment of dentinal hypersensitivity. For use in adults over the age of 21.

Contraindications: This product is contraindicated in patients with ulcerative gingivitis or stomatitis, or known sensitivity to silver or other heavy-metal ions. Patients with more than six affected sites, patients having had full mouth gingivectomies and patients showing abnormal skin sensitization in daily circumstances are recommended for exclusion.

Warnings: This product is intended for local application only. Not for ingestion. Protect the patient's eyes. Use caution to avoid contact with skin or clothing. In the event of exposure to eyes or skin, flush the area copiously with water and immediately seek medical consultation. This product yielded positive cytotoxicity in standard testing.

Precautions for Use:

- 1) Advantage Arrest does not normally stain enamel or burnished dentin. Advise patients that soft dentin or margins of composite restorations may be stained. Staining may be reversed by gentle polishing with tincture of iodine (weak iodine solution).
- 2) Advise patients that air-drying and product application can cause momentary transient pain to hypersensitive areas. Advantage Arrest has not been shown to cause pulpal necrosis even when soft dentin is treated.
- 3) Minimize product contact with gingiva and mucous membrane by using recommended amounts and careful application. Advantage Arrest may cause reversible short-term irritation. When applying Advantage Arrest to areas near the gingiva, apply petroleum jelly or cocoa butter and use cotton rolls to protect the gingival tissues. Alternatively, a rubber dam can be used to isolate the area.
- 4) If accidental contact occurs, thoroughly wash the area with water, saline solution or ~3% hydrogen peroxide. This includes contact with skin, clothes, floors and cabinets. Because Advantage Arrest is clear and thus

may be difficult to see, use caution to avoid transferring the material from gloved hands to other surfaces.

Precautions for Handling:

1. Storage Precautions

- 1) Store in original packaging in a cool, dark place.
- 2) Replace cap immediately after use.
- 3) Use as soon as dispensed.

2. Advantage Arrest will stain skin, clothes, counter tops, floors and instruments brown or black. Refer to the following for stain removal:

- 1) Skin; wash immediately with water, soap, ammonia or iodine tincture and then rinse thoroughly with water. Do not use excessive methods in an attempt to remove difficult stains from skin as the stains will eventually fade.
- 2) Clothing/Countertops/Floors/Instruments; use the same procedures as with stained skin. Difficult stains may be treated with sodium hypochlorite.

3. If Advantage Arrest is dispensed into a separate container, be sure to wash or thoroughly wipe the container clean immediately after use.

Adverse Reactions: Transient irritation of the gingiva has rarely been reported.

Dosage and Administration:

1. Isolate the affected area of the tooth with cotton rolls or protect the gingival tissue of the affected tooth with petroleum jelly. Alternatively, a rubber dam can be used to isolate the area.
2. Clean and dry the affected tooth surface.
3. For up to 5 treated sites per patient, dispense 1-2 drops of solution into a disposable dappen dish. Transfer material directly to the tooth surface with an applicator.
4. Air-dry.

If needed, one or two reapplications may be administered at intervals of one week.

How Supplied: Single 10 mL dropper-bottle containing 8 mL of product. Not sterile.

Storage: Do not freeze or expose to extreme heat. Keep in an air-tight container in a dark place.

Caution: Federal law restricts this device to sale by or on the order of a dentist or physician.

Distributed by:

Elevate Oral Care, LLC.
West Palm Beach, FL 33411
877-866-9113

© 2015 Elevate Oral Care

ELE137-0315

J.L. Castillo¹, S. Rivera¹, T. Aparicio²,
R. Lazo¹, T.-C. Aw³, L.L. Mancl⁴,
and P. Milgrom^{4*}

¹School of Dentistry, Universidad Peruana Cayetano Heredia, Lima, Peru; ²Private Practice, Cusco, Peru; ³Department of Restorative Dentistry, University of Washington, Seattle, USA; and ⁴Department of Dental Public Health Sciences, Box 357475, University of Washington, Seattle, WA 98195, USA; *corresponding author, dfrc@uw.edu

J Dent Res 90(2):203-208, 2011

ABSTRACT

Tooth sensitivity is a common clinical problem. This multi-center randomized clinical trial assessed the effectiveness and safety of topical diammine silver fluoride. From two sites (Lima and Cusco, Peru), 126 adults with at least one tooth sensitive to compressed air were randomly assigned to either the experimental treatment or sterile water, and pain was assessed by means of a 100-mm visual analogue scale at 24 hours and 7 days. The diammine silver fluoride reduced pain at 7 days at both sites. At the Lima site, the average change in pain scores between baseline and day 7 for the silver fluoride group was -35.8 (SD = 27.7) mm vs. 0.4 (SD = 16.2) mm for the control group ($P < 0.001$). In Cusco, the average change in pain scores for the silver fluoride group was -23.4 (SD = 21.0) mm and -5.5 (18.1) mm for the control group ($P = 0.002$). No tissue ulceration, white changes, or argyria was observed. A small number of participants in the silver fluoride group experienced a mild but transient increase in erythema in the gingiva near the tooth. No changes were observed in the Gingival Index. We concluded that diammine silver fluoride is a clinically effective and safe tooth desensitizer.

KEY WORDS: tooth sensitivity, silver diamine fluoride, diammine silver fluoride, silver diamine fluoride, fluorides, topical.

DOI: 10.1177/0022034510388516

Received August 13, 2010; Revised September 3, 2010;
Accepted September 3, 2010

© International & American Associations for Dental Research

The Short-term Effects of Diammine Silver Fluoride on Tooth Sensitivity: a Randomized Controlled Trial

INTRODUCTION

Tooth sensitivity to various stimuli, including cold air, has been explained by hydrodynamic changes within the dentinal tubules that activate intradental nerves (Markowitz and Pashley, 2008). Incidence is thought to be increasing. The etiology can be tooth wear, aggressive oral hygiene, and diet. Successful treatments physically block dentinal tubules (Arends *et al.*, 1997).

Sodium fluoride varnish and fluoride solutions and gels have been shown to reduce sensitivity (Thrash *et al.*, 1992; Ritter *et al.*, 2006). However, there is continuing interest in finding effective treatments. Nevertheless, recent studies have designs that are weak or statistically underpowered (Erdemir *et al.*, 2010; Jalali and Lindh, 2010).

The purpose of this study was to assess the clinical effectiveness and safety of topical diammine silver fluoride as a tooth desensitizer in adults.

METHODS

Design

This is a randomized clinical trial with two groups (Fig. 1). The study tested application of diammine silver fluoride in a single visit, because previous unpublished work had shown that a single application forms insoluble precipitates with calcium and phosphate that physically block dentinal tubules. The International Clinical Trials Registry number is NCT01063530.

Study Sites

The study was conducted in two sites, Lima and Cusco, Peru.

Participants

To be included, a participant must have at least one vital cuspid or premolar with a buccal cervical defect and clinical hypersensitivity in response to compressed air with a score ≥ 15 on a visual analogue scale (VAS) for pain. The individual will have had generally healthy gum tissue surrounding this tooth and no ulceration and no leukoplakia in this gingival tissue.

Candidates were excluded if they were using any type of tooth desensitizer, had received a fluoride varnish treatment within the preceding month, or were taking prescription medications, aspirin, or non-steroidal anti-inflammatory drugs; women who were pregnant were also excluded. Individuals using smokeless tobacco or chewing coca leaves were excluded. Individuals with known sensitivity to silver or other heavy-metal ions were excluded.




Figure 1. Flow chart for clinical trial sites combined.

Participants were recruited from the patient populations of Cayetano University School of Dentistry and the private dental practices of the investigators in Lima and Cusco between January and June, 2010, and were offered a small financial incentive for participation.

The Institutional Review Board of Universidad Peruana Cayetano Heredia approved the protocol, and the informed consent of all participants was obtained.

Treatment Conditions

Diammine silver fluoride [$\text{Ag}(\text{NH}_3)_2\text{F}$, CAS RN 33040–28–7, Saforide, Toyo Seiyaku Kasei Co. Ltd. Osaka, Japan] was used. It is clear and colorless, with a weak odor of ammonia. According to the manufacturer, the solution includes not less than 24.4 w/v% and not more than 26.8 w/v% of silver (Ag), not less than 5.0 w/v% and not more than 5.9 w/v% of fluorine (F). Diammine silver fluoride is also referred to as silver diammine fluoride, silver diammine fluoride, or silver fluoride.

Assignment to Conditions

Participants were randomly assigned to treatment with diammine silver fluoride or sterile water. The randomization was stratified on study site and baseline tooth sensitivity score (< 37 and ≥ 37) to a five-second blast of pressurized air at 2 cm distance from the tooth, and blocking was used to ensure that the

two groups would be balanced across the study period and within each stratum. The stratification at 37 was chosen from the literature (Ritter *et al.*, 2006). A pre-test of the VAS with 10 individuals confirmed the mean response in this range. Block sizes were equal to 2 or 4, and were chosen randomly with 2/3 and 1/3 probability, respectively. The assignments were generated by the project statistician, using the “sample” function of R statistical software (Version 2.7.1, The R Foundation for Statistical Computing, 2008). The assignments were recorded on slips of paper numbered consecutively within each stratum and then placed inside sealed envelopes sequentially numbered by stratum. The statistician retained the master list until all the data were analyzed. The clinician would open the envelope and apply the agent. The agents (active or control) were packaged in identical dark glass bottles labeled as A or B. The packaging was done at Cayetano University.

Clinical Procedure

The clinical procedure was that a disposable microbrush was dipped into a drop of the diammine silver fluoride or the control and then applied to the surface for 1 sec. Then the surface was gently air-dried and the procedure repeated.

Measures

Primary Outcome-Clinical

Reduction of pain (tooth sensitivity)—The teeth were isolated with gauze, and participants were asked to report tooth pain on a 100-mm visual analogue scale (VAS; Ritter *et al.*, 2006) before treatment and after treatment with a five-second blast of pressurized air at 2 cm distance from the tooth. The VAS was anchored with “no pain” and “intolerable pain”. The follow-up test was repeated at 24 hrs and 7 days later. A single person in each site conducted the assessment in Spanish. The scale was pre-tested to ensure that the descriptors were translated properly.

Safety

Damage to gingiva—Tissues were photographed before treatment to establish the normal baseline condition. A single examiner examined gingival tissues surrounding each treated tooth immediately after treatment, and at 24 hrs and 7 days later. The primary safety measure is erythema. It was assessed visually

Table 1. Tooth Sensitivity by Study Site and Condition

Study Site	Time	Condition		
Lima		Silver Fluoride (N = 37)	Control (N = 34)	
		Mean VAS (SD) [Range]	Mean VAS (SD) [Range]	P-value*
	Baseline	57.3 (26.7) [17, 99]	49.3 (19.3) [15, 84]	0.16
	24 hrs	28.2 (22.1) [2, 75]	52.1 (22.8) [16, 89]	
	Change from baseline	-29.1 (27.5) [-94, 10]	2.6 (15.3) [-44, 32]	< 0.0001
	7 days	21.5 (23.0) [1, 78]	49.9 (21.2) [9, 85]	
	Change from baseline	-35.8 (27.7) [-97, 12]	0.4 (16.2) [-38, 33]	< 0.0001
Cusco		Silver Fluoride (N = 26)	Control (N = 29)	
		Mean VAS (SD) [Range]	Mean VAS (SD) [Range]	P-value
	Baseline	51.7 (20.5) [22, 92]	51.6 (22.4) [16, 99]	0.98
	24 hrs	45.2 (24.1) [11, 87]	50.6 (22.0) [15, 95]	
	Change from baseline	-6.5 (13.1) [-34, 22]	-1.0 (11.7) [-37, 20]	0.11
	7 days	28.3 (21.8) [2, 94]	46.1 (24.4) [3, 92]	
	Change from baseline	-23.4 (21.0) [-56, 24]	-5.5 (18.1) [-77, 18]	0.0015

*Two-sample *t* test (unequal variances).

with the use of a standard dental light. Erythema (red changes) was rated on a 1 to 3 scale, where 1 is no redness, 2 is redness with bleeding on probing, and 3 is a severe change. The Gingival Index (Löe, 1967) was used to measure gingival inflammation in the mouth overall. White changes, ulceration, and staining were secondary measures. Changes were rated as present or absent. Examiners were trained to criteria using photographs and clinical cases. Intra- and inter-examiner reliability was established in 15 cases, and intraclass correlation was used to assess reliability. All intraclass correlations exceeded 0.8.

Data Analysis Plan

The data from the two sites were analyzed. To confirm reduction in pain, we calculated average difference scores between pre- and post-treatment VAS scores for each individual for each time-point (24 hrs and 7 days after treatment), and *t* tests were used to compare changes. The primary end point was at 7 days. Generalized estimating equations (GEE) linear regression was used in a secondary analysis to compare the reduction in pain across the 3 time-points, where the outcome is pain at the 3 time-points, the baseline pain is a covariate, and robust standard errors are used to account for multiple observations *per* participant and heteroscedasticity (Hardin and Hilbe, 2002). In addition, separate analyses of covariance were done at each time-point to compare the reduction in pain due to the active treatment between the two study sites, where the outcome is the pain at a particular time-point, baseline pain was entered as a covariate, and treatment and site, as well as a treatment-group-by-site interaction, were entered as factors.

We used Fisher’s Exact Test to assess whether there were more participants with erythema score > 1 in the silver fluoride group *vs.* the control group at 24 hrs and 7 days post-treatment. The primary end-point was assessed at 24 hrs. A *t* test assessed any differences in Gingival Index. Any white changes, ulceration, and staining (argyria) were reported.

Power Analysis

The data from the two sites were analyzed separately, and power is described below for the separate site analyses.

Reduction in tooth sensitivity—The primary end-point was assessed at 7 days post-treatment. In a similar desensitization study comparing fluoride varnishes (Ritter *et al.*, 2006), pain in response to air dropped from 36.9 (SD = 26.2) at baseline to 20.8 (SD = 4.3) at 2 wks post-treatment. We expected a similar or larger drop after 7 days with diammine silver fluoride, based on unpublished work from the University of Hong Kong, and little or no drop from the water. Thus, having 31 individuals in a group will allow for detection of effect size from 0.64 upwards, with an alpha of 0.05 and power of 0.8.

RESULTS

Participants

One hundred twenty-six adults (71 in Lima and 55 in Cusco) participated. About 378 candidates were screened between January and June 2010. The main reason (95%) for exclusion was lack of tooth sensitivity. The remainder were excluded because of the use of medications. No individuals were excluded because of tobacco use or coca. All of those eligible agreed to participate, but 10 were excluded because they failed to appear for the first visit. The proportion of women enrolled was 86% in Lima and 80% in Cusco. The average age of participants was 44 yrs and 43 yrs, respectively. There were no dropouts.

Participants and clinicians were blind to treatment assignment. Odor was not a threat to blinding, because the smell is not detectable clinically when such small quantities are used. Taste was not a threat in this study, because only minute amounts of material were applied and the tooth was air-dried after application.

Table 2. Numbers and Percentages of Participants with Erythema Score of 2 by Study Site and Condition

Study Site	Time	Condition		
Lima		Silver Fluoride (N = 37)	Control (N = 34)	
		n (%)	n (%)	P-value*
	Baseline	3 (8.1)	2 (5.9)	1.0
	24 hrs	3 (8.1)	2 (5.9)	1.0
	7 days	3 (8.1)	1 (2.9)	0.61
Cusco		Silver Fluoride (N = 26)	Control (N = 29)	
		n (%)	n (%)	P-value*
	Baseline	6 (23.1)	7 (24.1)	1.0
	24 hrs	10 (38.5)	2 (6.9)	0.0076
	7 days	3 (11.5)	3 (10.3)	1.0
Sites combined		Silver Fluoride (N = 63)	Control (N = 63)	
		n (%)	n (%)	P-value*
	Baseline	9 (14.3)	9 (14.3)	1.0
	24 hrs	13 (20.6)	4 (6.3)	0.035
	7 days	6 (9.5)	4 (6.3)	0.74

*Fisher's exact test

Clinical Effectiveness

The average pain scores before and after treatment, by site, are given in Table 1. At the Lima site, the silver fluoride group had slightly higher baseline scores (average = 57.3) than the control (average = 49.3; $P = 0.16$). At the Cusco site, the baseline scores were similar between the silver fluoride group (average = 51.7) and control (average = 51.6; $P = 0.98$). The primary study endpoint was the change from baseline to 7 days. In Lima, the average change in pain score between baseline and day 7 for the silver fluoride group was -35.8 (SD = 27.7) mm vs. 0.4 (SD = 16.2) for the controls ($P < 0.0001$). In Cusco, the average change in pain score between baseline and day 7 for the silver fluoride group was -23.4 (SD = 21.0) mm vs. -5.5 (SD = 18.1) mm ($P = 0.0015$) for water.

Comparison of tooth sensitivity at 24 hrs and 7 days between study groups by analysis covariance, adjusted for the baseline sensitivity level, gave similar results.

There was no significant three-way interaction among study site, time, and study group (GEE linear regression; $P = 0.20$), but all two-way interactions were significant: study site by time ($P = 0.043$), study site by study group ($P = 0.0006$), and study group by time ($P = 0.0076$). Hence, an analysis of time effect was done separately by study site. In Lima, there was no significant time-by-study-group interaction ($P = 0.21$). The overall study group difference in tooth sensitivity (over both time-points), adjusted for baseline sensitivity, was 29.9 ($P < 0.001$). The overall difference in sensitivity between 24 hrs and 7 days was 4.5 ($P = 0.014$). In Cusco, there was a significant study-group-by-time interaction ($P = 0.015$), so the overall study group difference is not reported. The differences in sensitivity between 24 hrs and 7 days were 16.9 ($P = 0.005$) for silver fluoride and 4.5 ($P = 0.097$) in the control group, respectively.

Safety

The number and percent of participants with a erythema score of 2 for the gingival tissue of the tooth treated for each treatment condition by site and time are given in Table 2. Scores were low; no individual had score 3, severe erythema, either before or after the application of silver fluoride. There was no difference in the proportion of participants with erythema score > 1 between the silver fluoride group and the placebo (Fisher's Exact Test, $P = 1.0$) at any time-point in the Lima population. There was a small but significant increase in the proportion of participants at the Cusco site who experienced an erythema score > 1 at 24 hrs ($P = 0.0076$). There was no difference in the proportion of participants with an erythema score > 1 between the groups in Cusco after 7 days ($P = 1.0$). No white or dark changes were noted in gingiva in any participant at any time in any condition at either site. An independent examiner, who was blind to treatment condition and time, examined the photographs and confirmed this lack of change.

The Gingival Index scores for each treatment condition and site are listed in Table 3. The mean (SD) Gingival Index scores for the mouth for treatment and control groups at baseline were: (Lima) silver fluoride, 0.29 (0.24), control 0.33 (0.35) ($P = 0.59$); and (Cusco) silver fluoride, 0.47 (0.24), control 0.38 (0.27) ($P = 0.19$). At 7 days, the mean (SD) changes in GI scores were: (Lima) silver fluoride, -0.02 (0.09), control 0.03 (0.13) ($P = 0.076$); and (Cusco) silver fluoride, -0.16 (0.27), control -0.03 (0.09) ($P = 0.023$). Similar results were observed after 24 hrs.

Photographs of the teeth suggest that the silver fluoride did not stain most exposed root surfaces (see Fig. 2 for an example). This result was found only when surfaces had untreated decay.

Table 3. Overall Gingival Index Score by Study Site and Condition

Study Site	Time	Condition		
		Silver Fluoride (N = 37)	Control (N = 34)	
		Mean (SD) [Range]	Mean (SD) [Range]	P-value*
Lima	Baseline	0.29 (0.24) [0.0, 1.2]	0.33 (0.35) [0.0, 1.5]	0.59
	24 hrs	0.28 (0.24) [0.0, 1.2]	0.35 (0.36) [0.0, 1.7]	
	Change from baseline	-0.01 (0.05) [-0.2, 0.1]	0.02 (0.07) [-0.2, 0.2]	0.076
	7 days	0.27 (0.23) [0.0, 1.2]	0.36 (0.39) [0.1, 1.8]	
	Change from baseline	-0.02 (0.09) [0.2, 0]	0.03 (0.13) [-0.5, 0.3]	0.076
Cusco	Baseline	0.47 (0.24) [0.1, 0.9]	0.38 (0.27) [0.0, 1.2]	0.19
	24 hrs	0.36 (0.21) [0.1, 0.8]	0.36 (0.24) [0.0, 1.2]	
	Change from baseline	-0.11 (0.16) [-0.6, 0.1]	-0.02 (0.12) [-0.3, 0.3]	0.020
	7 days	0.31 (0.19) [0.0, 0.8]	0.35 (0.26) [0.1, 1.2]	
	Change from baseline	-0.16 (0.27) [-0.8, 0.7]	-0.03 (0.09) [-0.3, 0.2]	0.023
Sites Combined	Baseline	0.36 (0.26) [0.0, 1.2]	0.35 (0.32) [0.0, 1.5]	0.72
	24 hrs	0.31 (0.23) [0.0, 1.2]	0.35 (0.31) [0.0, 1.7]	
	Change from baseline	-0.05 (0.12) [-0.6, 0.1]	0.00 (0.10) [-0.3, 0.3]	0.0023
	7 days	0.28 (0.22) [0.0, 1.2]	0.35 (0.33) [0.1, 1.8]	
	Change from baseline	-0.08 (0.20) [-0.8, 0.7]	0.00 (0.12) [-0.5, 0.3]	0.0028

*Two-sample test (unequal variances).

**Analysis of covariance, adjusted for study site, with heteroscedasticity-consistent standard errors.

DISCUSSION

In a population with teeth sensitive to air, this trial demonstrated that a topical solution of diammine silver fluoride was more effective than a placebo in reducing tooth pain. Reductions grew larger between 24 hrs and 7 days post-treatment. The study was conducted in two sites by different investigators to increase generalizability and had sufficient statistical power to detect clinically meaningful differences in pain. The study involved many more individuals than the typical study (Ritter *et al.*, 2006).

The results, however, are consistent with those from similar studies of other desensitizers, such as self-administered 0.717% fluoride solution (Thrash *et al.*, 1992) or fluoride varnish (Ritter *et al.*, 2006). In the fluoride solution study, the authors concluded that two one-minute applications reduced sensitivity to cold. Participants in the varnish study experienced a pain reduction in response to ice, but not to air, at 2 wks. The current study reported significant pain reductions in response to air in 24 hrs that were maintained at 7 days. The magnitude of reduction was considerably greater than in the other studies. The current study did not use ice as a stimulus.

There were no unintended effects on the gingiva, and any inflammation resulting from the treatment was minor and transient. No staining of the gingival tissues was observed.



Figure 2. Root caries at baseline (left panel), 24 hrs after treatment (middle panel), and 7 days after treatment with diammine silver fluoride (right panel).

Staining of teeth was found only when surfaces had untreated decay. The staining of carious dentin can be minimized by the application of potassium iodide solution after treatment without reducing the effect (Knight *et al.*, 2006).

Diammine silver fluoride has been shown to arrest caries in animal models (Tanzer *et al.*, 2010) and to be more effective than sodium fluoride varnish in human trials (Chu *et al.*, 2002; Llodra *et al.*, 2005; Rosenblatt *et al.*, 2009; Tan *et al.*, 2010). It did not cause abscesses in teeth with open cavities that were treated. The mechanism of action for caries arrest may be antimicrobial (Knight *et al.*, 2009). Studies have also shown that diammine silver fluoride is free of adverse effects (Chu *et al.*, 2002; Llodra *et al.*, 2005; Tan *et al.*, 2010). This suggests that diammine fluoride may be particularly effective in individuals in whom sensitivity is associated with demineralization and caries.

Diammine silver fluoride has been demonstrated to be a clinically effective and safe tooth desensitizer after 24 hrs and 7 days. Clinical trials are warranted to examine effectiveness over a longer period of time and in comparison with other agents.

ACKNOWLEDGMENTS

The authors acknowledge the contributions of Silvia Navarro in recruitment of participants. ADP Silver Dental Arrest, LLC, Redmond, OR, USA, was the study sponsor.

REFERENCES

- Arends J, Düschner H, Ruben JL (1997). Penetration of varnishes into demineralized root dentin *in vitro*. *Caries Res* 31:201-205.
- Chu CH, Lo EC, Lin HC (2002). Effectiveness of silver diamine fluoride and sodium fluoride varnish in arresting dentin caries in Chinese pre-school children. *J Dent Res* 81:767-770.
- Erdemir U, Yildiz E, Kilic I, Yucel T, Ozel S (2010). The efficacy of three desensitizing agents used to treat dentin hypersensitivity. *J Am Dent Assoc* 141:285-296.
- Hardin JW, Hilbe JM (2002). Generalized estimating equations. New York, NY: Chapman & Hall/CRC Press.
- Jalali Y, Lindh L (2010). A randomized prospective clinical evaluation of two desensitizing agents on cervical dentine sensitivity. A pilot study. *Swed Dent J* 34:79-86.
- Knight GM, McIntyre JM, Craig GG, Mulyani (2006). Ion uptake into demineralized dentine from glass ionomer cement following pretreatment with silver fluoride and potassium iodide. *Aust Dent J* 51: 237-241.
- Knight GM, McIntyre JM, Craig GG, Mulyani, Zilm PS, Gully NJ (2009). Inability to form a biofilm of *Streptococcus mutans* on silver fluoride- and potassium iodide-treated demineralized dentin. *Quintessence Int* 40:155-161.
- Llodra JC, Rodriguez A, Ferrer B, Menardia V, Ramos T, Morato M (2005). Efficacy of silver diamine fluoride for caries reduction in primary teeth and permanent first molars of schoolchildren: 36-month clinical trial. *J Dent Res* 84:721-724.
- Lõe H (1967). The Gingival Index, the Plaque Index, and the Retention Index. *J Periodontol* 38(Suppl):610-616.
- Markowitz K, Pashley DH (2008). Discovering new treatments for sensitive teeth: the long path from biology to therapy. *J Oral Rehabil* 35: 300-315.
- Ritter AV, Dias WdeL, Miguez P, Caplan DJ, Swift EJ Jr (2006). Treating cervical dentin hypersensitivity with fluoride varnish: a randomized clinical study. *J Am Dent Assoc* 137:1013-1020.
- Rosenblatt A, Stamford TC, Niederman R (2009). Silver diamine fluoride: a caries "silver bullet". *J Dent Res* 88:116-125.
- Tan HP, Lo EC, Dyson JE, Luo Y, Corbet EF (2010). A randomized trial on root caries prevention in elders. *J Dent Res* 89(10):1086-1090. Epub 2010 Jul 29.
- Tanzer J, Thompson A, Milgrom P, Shirtcliff M (2010). Diammine silver fluoride arrestment of caries associated with anti-microbial action (abstract). *J Dent Res* 89(Spec Iss B):2082. URL: <http://iadr.confex.com/iadr/2010barce/webprogram/Paper131877.html> (URL accessed 09/07/2010).
- Thrash WJ, Jones DL, Dodds WJ (1992). Effect of a fluoride solution on dentinal hypersensitivity. *Am J Dent* 5:299-302.

UCSF Protocol for Caries Arrest Using Silver Diamine Fluoride: Rationale, Indications and Consent

Jeremy A. Horst, DDS, PhD; Hellene Ellenikiotis, DDS; and Peter L. Milgrom, DDS

ABSTRACT The Food and Drug Administration recently cleared silver diamine fluoride for reducing tooth sensitivity. Clinical trials document arrest and prevention of dental caries by silver diamine fluoride. This off-label use is now permissible and appropriate under U.S. law. A CDT code was approved for caries arresting medicaments for 2016 to facilitate documentation and billing. We present a systematic review, clinical indications, clinical protocol and consent procedure to guide application for caries arrest treatment.

AUTHORS

Jeremy A. Horst, DDS, PhD, is a fellow at the University of California, San Francisco, School of Dentistry studying the bacteria that cause cavities, a pediatric dentist at Alameda Pediatric Dentistry and co-founder and CSO at OraViz.

Conflict of Interest

Disclosure: Dr. Horst is co-founder and CSO at OraViz.

Hellene Ellenikiotis, DDS, is a resident in the University of California, San Francisco, general practice residency and a recent graduate of the University of California, San Francisco School of Dentistry.

Conflict of Interest

Disclosure: None reported.

Peter M. Milgrom, DDS, is a professor of dental public health sciences and pediatric dentistry and director of the Northwest Center to Reduce Oral Health Disparities at the University of Washington in Seattle.

Conflict of Interest

Disclosure: Dr. Milgrom is a principal in ADP Silver Dental Arrest LLC, which licenses permission to market Advantage Arrest to Elevate Oral Care LLC.

Until now, no option for the treatment of dental caries in the U.S. besides restorative dentistry has shown substantial efficacy.¹ Silver diamine fluoride is an inexpensive topical medicament used extensively in other countries to treat dental caries across the age spectrum. No other intervention approaches the ease of application and efficacy. Multiple randomized clinical trials — with hundreds of patients each — support its use for caries treatment, thus substantiating an intervention that addresses an unmet need in American dentistry. In August 2014, the Food and Drug Administration (FDA) cleared the first silver diamine fluoride product for market, and as of April 2015, that product is available.

Since its approval in Japan more than 80 years ago,² more than 2 million containers have been sold. The silver acts as an antimicrobial, the fluoride promotes remineralization and the ammonia stabilizes high concentrations in solution.³

Because silver diamine fluoride is new to American dentistry and dental education, there is a need for a standardized guideline, protocol and consent. The University of California, San Francisco, School of Dentistry paradigm shift committee assembled a subcommittee with the following goals:

- Use available evidence to develop a list of clinical indications.
- Define a protocol that maximized safety and efficacy and minimized inadvertent staining of clinical facilities.

- Build an informed consent document at the eighth-grade reading level.

We conducted a systematic review, inquired of authors of published clinical and in vitro studies about details and considerations in their protocols and consulted experts in cariology and materials chemistry where evidence was lacking. The work of this committee resulted in the adoption of silver diamine fluoride use in the UCSF student clinics.

Methods

A literature review was designed by a medical librarian to search PubMed and the International Association of Dental Research abstract archive with the following search terms: “33040-28-7” OR “1Z00ZK3E66” OR “silver diamine fluoride” OR “silver fluoride” OR “silver diamine fluoride” OR “diammine silver fluoride” OR “ammonical silver fluoride” OR “ammoniacal silver fluoride”. Differences in nomenclature have led to confusion around this material. Another review was completed with the terms “dental” OR “caries” AND “silver nitrate” AND “clinical.”

Material

Silver diamine fluoride (38% w/v $\text{Ag}(\text{NH}_3)_2\text{F}$, 30% w/w) is a colorless topical agent comprised of 24.4-28.8% (w/v) silver and 5.0-5.9% fluoride at pH 10,⁴ and marketed as Advantage Arrest by Elevate Oral Care LLC (West Palm Beach, Fla.). Other companies may market silver diamine fluoride in the future following determination of substantial equivalence and FDA clearance.

Mechanisms

Silver diamine fluoride is used for caries arrest and treatment of dentin hypersensitivity. In the treatment of exposed sensitive dentin surfaces,

topical application results in the development of a squamous layer on the exposed dentin, partially plugging the dentinal tubules.⁵ High concentration aqueous silver has been long known to form this protective layer.⁶ Decreased sensitivity in treated patients^{7,8} is consistent with the hydrodynamic theory of dentin hypersensitivity.⁹

Dental caries is a complex progression involving dietary sugars, bacterial metabolism, demineralization and organic degradation. The collagenous organic matrix is

Silver diamine fluoride outperforms other anticaries medicaments in killing cariogenic bacteria in dentinal tubules.

exposed once a dentin surface is demineralized and destroyed by native and bacterial proteases to enable a lesion to enlarge.¹⁰ Upon application of silver diamine fluoride to a decayed surface, the squamous layer of silver protein conjugates forms, increasing resistance to acid dissolution and enzymatic digestion.¹¹ Hydroxyapatite and fluorapatite form on the exposed organic matrix, along with the presence of silver chloride and metallic silver.⁵ The treated lesion increases in mineral density and hardness while the lesion depth decreases.⁵ Meanwhile, silver diamine fluoride specifically inhibits the proteins that break down the exposed dentin organic matrix: matrix metalloproteinases,¹¹ cathepsins¹² and bacterial collagenases.⁵ Silver

ions act directly against bacteria in lesions by breaking membranes, denaturing proteins and inhibiting DNA replication.^{13,14} Ionic silver deactivates nearly any macromolecule. Silver diamine fluoride outperforms other anticaries medicaments in killing cariogenic bacteria in dentinal tubules.¹⁵

Silver and fluoride ions penetrate ~25 microns into enamel¹⁶ and 50-200 microns into dentin.¹⁷ Fluoride promotes remineralization, and silver is available for antimicrobial action upon release by re-acidification.¹⁸ Silver diamine fluoride arrested lesions are 150 microns thick.¹⁹

Artificial lesions treated with silver diamine fluoride are resistant to biofilm formation and further cavity formation, presumably due to remnant ionic silver.^{20,21} More silver and fluoride is deposited in demineralized than nondemineralized dentin. Correspondingly, treated demineralized dentin is more resistant to caries bacteria than treated sound dentin.²² When bacteria killed by silver ions are added to living bacteria, the silver is re-activated so that effectively the dead bacteria kill the living bacteria in a “zombie effect.”²³ This reservoir effect helps explain why silver deposited on bacteria and dentin proteins within a cavity has sustained antimicrobial effects.

Clinical Evidence

Silver Nitrate Plus Fluoride Varnish

Before the FDA cleared silver diamine fluoride, some U.S. dentists sequentially applied silver nitrate then fluoride varnish to dentinal decay as the only available noninvasive option for caries treatment. Duffin rediscovered silver nitrate from the early literature,²⁴ which had been lost

to modern cariology. Surprisingly, there is no mention of silver nitrate in either of the American Dental Association Council on Scientific Affairs reports on *Nonfluoride Caries-Preventive Agents*²⁵ or *Managing Xerostomia and Salivary Gland Hypofunction*,²⁶ and it is not part of standard dental school curricula. Case series of carious lesions arrested by silver nitrate date to the 1800s. For example, in 1891, 87 of 142 treated lesions were arrested.²⁷ Percy Howe, DDS, then director of the Forsyth Institute in Boston, added ammonia to silver nitrate, making it more stable and effective as an antimicrobial for application to any infected tooth structure from early cavitated lesions to infected root canals.²⁸ Duffin added the application of fluoride varnish following silver nitrate, simulating silver diamine fluoride. While his clinic doubled in patients, cases needing general anesthesia disappeared. His review of randomly selected charts showed only seven of 578 treated lesions progressed within two and a half years to the point that extractions were needed.²⁴ Thus, with the exception of Duffin's and one other report, attention to silver nitrate largely disappeared by the 1950s. The lore is that use and teaching of this intervention were lost with the introduction of effective local anesthetic to enable painless restorations and fluoride for caries prevention. Because no high-quality clinical trials have been performed, we did not include the silver nitrate plus fluoride varnish regimen in our recommendation.

Silver Diamine Fluoride

We found nine published randomized clinical trials evaluating silver diamine fluoride for caries arrest and/or prevention of at least one year in duration. These

studies each involved hundreds of children aged 3 to 9 or adults aged 60 to 89 (**FIGURES 1** and **2**). Most participants had low (< 0.3 ppm) fluoride in the environmental water and reported using fluoride toothpaste (e.g., 73 percent).²⁹ Silver diamine fluoride was applied with cotton isolation. Lesions were detected with mirror and explorer only. All studies were registered and met the Consolidated Standards of Reporting Trials requirements. Clinical cases and studies not meeting these criteria can be found elsewhere.³⁰

When stannous fluoride was used to activate color change, a break in the black color within a lesion at six months was highly sensitive and specific for active caries.

Caries arrest increased dramatically after reapplication from one year posttreatment³¹⁻³³ to one and a half years,^{31,34} and increasingly to two to three years (**FIGURE 1**).^{29,31,35} Single application without repeat lost effect over time in the elderly.³² Twice per year application resulted in more arrest than once per year.^{31,35} Twelve percent silver diamine fluoride was markedly less effective.³²

Darkening of the entire lesion indicated success at follow-up and is suggested to facilitate diagnosis of caries arrest status by nondentists. A longitudinal study reported that color activation of silver diamine fluoride with 10% stannous fluoride resulted in less first molar caries.³⁶ Tea extract was used in one group to activate color change for improved follow-up diagnosis; no differences

in arrest were seen.³² Indeed, when stannous fluoride was used to activate color change, a break in the black color within a lesion at six months was highly sensitive and specific for active caries.³⁷

Silver diamine fluoride greatly outperformed fluoride varnish for caries arrest²⁹ and was equivalent or better than glass ionomer cement (GIC) (**FIGURE 1**).^{31,33} The addition of semiannual intensive oral health education with the application of silver diamine fluoride in the elderly increased the arrest of root caries (**FIGURE 1**).³⁸

Caries Prevention

When silver diamine fluoride was applied only to carious lesions, impressive prevention was seen for other tooth surfaces.^{29,35} Fluoride-releasing GIC can have this effect but it is limited to surfaces adjacent to the treated surface and of short duration. Direct application to healthy surfaces in children also helps prevent caries (**FIGURE 2**).^{29,35,39} Two studies show great difference in the level of prevention in the elderly;^{38,40} the difference is hard to reconcile. As seen for arrest, prevention is less after one year without repeat application.⁴¹

Annual application of silver diamine fluoride prevented many more carious lesions than four-times-per-year fluoride varnish in both children²⁹ and the elderly.⁴⁰ Prevention was roughly equivalent to twice-per-year varnish in one study (**FIGURE 2**).³⁹ The addition of semiannual intensive oral health education in a study of the elderly increased prevention.³⁸ Although many fell out, GIC or resin sealants outperformed silver diamine fluoride in preventing caries in the first molars of children,^{39,41} though the cost was ~20 times more.

Ongoing Trials

Unpublished reports of clinical studies unanimously confirm better caries arrest and/or prevention by silver diamine fluoride over control or other materials. A one-year report of a study of the elderly demonstrated that the addition of a saturated solution of potassium iodide (SSKI) to decrease discoloration did not significantly alter caries arrest or prevention.⁴² This was confirmed in the two-year examinations (personal communication, Edward Lo). A one-year report of a study in children showed that the application once per week for three consecutive weeks, once per year, was more effective than that of single annual application.⁴³ Other studies have recently begun to evaluate the ability of silver diamine fluoride to arrest interproximal carious lesions, to compare the relative efficacy of silver diamine fluoride to the combination of silver nitrate plus fluoride varnish and to compare the effects on populations with or without access to fluoridated water. Final reports from these studies will follow in the coming years.

Recommendations From the Literature on Clinical Efficacy

These studies show that 38% silver diamine fluoride is effective and efficient in arresting and preventing carious lesions. Application only to lesions appears to be similarly effective in preventing cavities in other teeth and surfaces as applying directly. Single application appears insufficient for sustained effects, while annual re-application results in remarkable success, and even greater effects with semi-annual application. From these data, we recommend twice-per-year application, only to carious lesions without excavation, for at least the first two years.




FIGURE 1. Graphic summary of randomized controlled trials demonstrating caries arrest after topical treatment with 38% silver diamine fluoride (SDF). Studies are arranged vertically by frequency of silver diamine fluoride application. Caries arrest is defined as the fraction of initially active carious lesions that became inactive and firm to a dental explorer. SDF (38% unless noted otherwise); q6mon, every six months; q1year, every year; q3mon, every three months; GIC, glass ionomer cement; NaF, 5% sodium fluoride varnish; + OHI q6mon, SDF every year and oral hygiene instructions every six months.

For any patient with active caries, we recommend considering replacement of fluoride varnish as the primary means to prevent new lesions, with application of silver diamine fluoride to the active lesions only. For patients without access

to both sealants and monitoring, silver diamine fluoride is the agent of choice for prevention of caries in permanent molars — particularly as there is no margin to leak and thereby facilitate deep caries and it does not stain sound enamel.




FIGURE 2. Graphic summary of randomized controlled trials demonstrating caries prevention after topical treatment of carious lesions with 38% silver diamine fluoride. Prevented caries is defined as the fraction of new carious lesions in treatment groups as compared to those in the placebo or no treatment control group. Chlorhex, 1% chlorhexidine varnish.

Longer studies are needed to determine whether caries arrest and prevention can be maintained with decreased application after two to three years, and whether more frequent use would enhance efficacy. Traditional or

nontraditional restorative approaches, such as the atraumatic restorative technique (ART)⁴⁴ and Hall crowns,⁴⁵ should be performed as dictated by the response of the patient, disease progression and the nature of individual lesions.

Safety

Maximum Dose and Safety Margin

The margin of safety for dosing is of paramount concern. In gaining clearance from the FDA, female and male rat and mouse studies were conducted to determine the lethal dose (LD50) of silver diamine fluoride by oral and subcutaneous administration. Average LD50 by oral administration was 520 mg/kg and by subcutaneous administration was 380 mg/kg. The subcutaneous route is taken here as a worst-case scenario. One drop (25 μ L) is ample material to treat five teeth and contains 9.5 mg silver diamine fluoride. Assuming the smallest child with caries would be in the range of 10 kg, the dose would be 0.95 mg/kg child. Thus, the relative safety margin of using an entire drop on a 10 kg child is 380 mg/kg LD50/0.95 mg/kg dose = four-hundredfold safety margin. The actual dose is likely to be much smaller, for example 2.37 mg total for three teeth was the largest dose measured in six patients.⁴⁶ The most frequent application monitored in a clinical trial was weekly for three weeks, annually.⁴³ Thus, we set our recommended limit as one drop (25 μ L) per 10 kg per treatment visit, with weekly intervals at most. This dose is commensurate with the Environmental Protection Agency's (EPA) allowable short-term exposure of 1.142 mg silver per liter of drinking water for one to 10 days (Agency for Toxic Substances and Disease Registry, ATSDR, 1990).

Cumulative exposure from lower-level acute or chronic silver intake has no real physiologic disease importance, but the bluing of skin in argyria should obviously be avoided. The EPA set the lifetime exposure conservatively at 1 gm to safely avoid argyria. The highest applied dose for three teeth measured in the pharmacokinetic study, 2.37 mg, would enable > 400 applications.⁴⁶ Silver

nitrate (typically a 25% solution) has been used for more than 100 years in the U.S. without incident, including acceptance by the ADA, and in other countries for arresting dental caries.³

Adverse Effects

Not a single adverse event has been reported to the Japanese authorities since they approved silver diamine fluoride (Saforide, Toyo Seiyaku Kasei Co. Ltd., Osaka, Japan) more than 80 years ago.⁴⁷ The manufacturer estimates that more than 2 million multi-use containers have been sold, including > 41,000 units in each of the last three reporting years.

In the nine randomized clinical trials in which silver diamine fluoride was applied to multiple teeth to arrest or prevent dental caries, the only side effect noted was for three of 1,493 children or elderly patients monitored for one to three years who experienced “a small, mildly painful white lesion in the mucosa, which disappeared at 48 [hours] without treatment.”^{29,31-33,35,38,40,41,48} The occurrence of reversible localized changes to the oral mucosa was predicted in the first reports of longitudinal studies.⁴⁹ No adverse pulpal response was observed.

Gingival responses have been minimal. In a pharmacokinetic study of silver diamine fluoride application to three teeth in each of six 48- to 82-year-olds, no erythema, bleeding, white changes, ulceration or pigmentation was found after 24 hours. Serum fluoride hardly went up from baseline, while serum silver increased about tenfold and stayed high past the four hours of measurement.⁴⁶ In a two-site hypersensitivity trial of 126 patients in Peru, at baseline 9 percent of patients presented redness scores of 2 (1 being normal, 2 being mild to moderate redness and 3 being severe); and after one day, 13 percent in silver diamine fluoride treated patients versus 4 percent

in controls. All redness was gone at seven days. Meanwhile, gingival index improved slightly in silver diamine fluoride treated patients.⁷ Nonetheless, gingival contact should be minimized. In our experience, it has been adequate to coat the nearby gingiva with petroleum jelly, use the smallest available microsponge and dab the side of the dappen dish to remove excess liquid before application.

Concerns for fluoride safety are most relevant to chronic exposure,⁵⁰ whereas this is an acute exposure. Chronically high systemic fluoride results in dental

At least for children, many parents have seen the color changes as a positive indication that the treatment was effective.

fluorosis. The ubiquitous use of fluoride-based gas in general anesthetics has shown that the first acute response is transient renal holding, and is rare.⁵¹ Concerns have been raised about poorly controlled silver diamine fluoride concentrations⁵² and fluorosis appearing in treated rats.⁵³ However, silver and fluoride levels are closely monitored for the U.S. product, and the Health Department of Western Australia conducted a study that found no evidence of fluorosis resulting from long-term proper use of silver diamine fluoride.⁵⁴ Therefore, we have concluded that the development of fluorosis after application of the U.S.-approved product is not a clinically significant risk.

Silver allergy is a contraindication. Relative contraindications include any significant desquamative gingivitis or

mucositis that disrupts the protective barrier formed by stratified squamous epithelium. Increased absorption and pain would be expected with contact. Heightened caution and use of a protective gingival coating may suffice.

A saturated solution of potassium iodide (SSKI) is contraindicated in pregnant women and during the first six months of breastfeeding because of the concern of overloading the developing thyroid with iodide; thyroid specialists suggested a pregnancy test prior to use in women of childbearing age uncertain of their status.

Nonmedical Side Effects

Silver diamine fluoride darkens carious lesions. At least for children, many parents have seen the color changes as a positive indication that the treatment was effective.²⁹ Application of an SSKI immediately following silver diamine fluoride treatment is thought to decrease staining (patent US6461161). This is an off-label use; potassium iodide is approved as an over-the-counter drug to facilitate mucus release to breathe more easily with chronic lung problems and to protect the thyroid from radioactive iodine in radiation emergencies. In our clinical experience, SSKI helps but does not dramatically effect stain; arrested lesions normally darken. Most stain remains at the dentin-enamel or cementum-enamel junction. However, SSKI maintains resistance to biofilm formation or activity in laboratory studies.²⁰ Also, SSKI maintained caries arrest efficacy in the early results of an ongoing clinical trial.⁴² Meanwhile, silver diamine fluoride-treated lesions can also be covered with GIC or composite (see below for discussion on bonding).

Patients note a transient metallic or bitter taste. In our experience, with judicious use, the taste and texture

response is more favorable than the response to fluoride varnish.

Even a small amount of silver diamine fluoride can cause a “temporary tattoo” to the skin (on the patient or provider), like a silver nitrate stain or henna tattoo, and does no harm. Stain on the skin resolves with the natural exfoliation of skin in two to 14 days. Universal precautions prevent most exposures. Long-term mucosal stain, local argyria akin to an amalgam tattoo, has been observed when applying silver nitrate to intraoral wounds; we anticipate similar stains with submucosal exposure to silver diamine fluoride.

Silver diamine fluoride stains clinic surfaces and clothes. The stain does not come out once it sets. Spills should be cleaned up immediately with copious water, ethanol or bleach. High pH solvents such as ammonia may be more successful. Secondary containers and plastic liners for surfaces are adequate preventives.

Effects on Bonding

Using a contemporary bonding system, silver diamine fluoride had no effect on composite bonding to noncarious dentin using either self-etch or full-etch systems.⁵⁵ In one study, simply rinsing after silver diamine fluoride application avoided a 50 percent decrease in bond strength for GIC.⁵⁶ In another study, increased dentin bond strength to GIC was observed.⁵⁷ Silver diamine fluoride decreased dentin bonding strength of resin-based crown cement by approximately one-third.⁵⁸ Thus, rinsing will suffice for direct restorations, while excavation of the silver diamine fluoride-treated superficial dentin is appropriate for cementing crowns.

Indications

Countless patients would benefit from conservative treatment of nonsymptomatic active carious lesions. We discuss the following indications.

First, extreme caries risk is defined as patients with salivary dysfunction, usually secondary to cancer treatment, Sjogren’s syndrome, polypharmacy, aging or methamphetamine abuse. For these patients, frequent prevention visits and traditional restorations fail to stop disease progression. Similar disease recurrence occurs in severe early childhood caries.

Second, some patients cannot tolerate standard treatment for medical or psychological reasons. These include the preoperative child, the frail elderly, those with severe cognitive or physical

Countless patients would benefit from conservative treatment of nonsymptomatic active carious lesions.

disabilities and those with dental phobias. Various forms of immunocompromise mean that these same patients have a much higher risk of systemic infection arising from untreated dental caries. Many only receive restorative care with general anesthesia or sedation and others are not good candidates for general anesthesia due to frailty or another medical complexity. The Centers for Disease Control and Prevention (CDC) estimates 1.4 million people in the U.S. live in nursing homes and 1.2 million live in hospice.⁵⁹ These individuals tend to have medical, behavioral, physical and financial limitations that beg a reasonable option.

Third, some patients have more lesions than can be treated in one visit, such that new lesions arise or existing lesions become symptomatic while

awaiting completion of treatment. This is particularly relevant to the dental school setting where treatment is slow. American dentistry has been desperately lacking an efficient instrument to be used at the diagnostic visit to provide a step toward controlling the disease.

Fourth, some lesions are just difficult to treat. Recurrent caries at a crown margin, root caries in a furcation or the occlusal of a partially erupted wisdom tooth pose a challenge to access, isolation and cleansability necessary for restorative success.

Following the above considerations, we developed four indications for treatment of dental caries with silver diamine fluoride:

1. Extreme caries risk (xerostomia or severe early childhood caries).
2. Treatment challenged by behavioral or medical management.
3. Patients with carious lesions that may not all be treated in one visit.
4. Difficult to treat dental carious lesions.

Finally, these indications are for our school clinics. They do not address access to care. The U.S. Department of Health and Human Services estimates 108 million Americans are without dental insurance, and there are 4,230 shortage areas with 49 million people without access to a dental health professional.⁶⁰ Unlike fillings, failure of silver diamine fluoride treatment does not appear to create an environment that promotes caries, and thus needs to be monitored. Thus, a final important indication is:

5. Patients without access to dental care.

Clinical Application

We considered practical strategies to maximize safety and effectiveness in the design of a clinical protocol for the UCSF dental clinics (**FIGURE 3**).

The key factor is repeat application

Silver Diamine Fluoride (SDF) UCSF Protocol for Arresting Dental Carious Lesions or Treating Tooth Sensitivity

Material: Advantage Silver Arrest (38% SDF, purified water) from Elevate Oral Care.
Shelf life: three years unopened. Do not refrigerate. Avoid freezing or extreme heat.

Indications:

1. Extreme caries risk (xerostomia or severe early childhood caries).
2. Treatment challenged by behavioral or medical management.
3. Patients with carious lesions that may not all be treated in one visit.
4. Difficult to treat dental carious lesions.
5. Patients without access to dental care.

Maximum dose: 25 µL (1 drop) / 10kg per treatment visit.

SDF Contraindication: Silver allergy.

SDF Relative Contraindications: Ulcerative gingivitis, stomatitis.

SSKI Contraindications: Pregnancy, breastfeeding.

Considerations:

- Decayed dentin will darken as the caries lesions arrest. Most will be dark brown or black.
- SDF can stain the skin, which will clear in two to three weeks without treatment.
- SDF can permanently stain operatory surfaces and clothes.
- A control restoration (e.g., GI via ART or other material) may be considered after SDF treatment.
- Saturated solution of potassium iodide (SSKI, Lugol's Solution, various sources) can be used after SDF to decrease color changes.
- Re-application is usually recommended, biannually until the cavity is restored or arrested or the tooth exfoliates.

Procedure:

1. Plastic-lined cover for counter, plastic-lined bib for patient.
2. Standard personal protective equipment (PPE) for provider and patient.
3. One drop of SDF into the deep end of a plastic dappen dish (also obtain one drop of SSKI in a separate dappen dish if selected).
4. Remove bulk saliva with saliva ejector.
5. Isolate tongue and cheek from affected teeth with 2-inch by 2-inch gauze or cotton rolls.
6. If near the gingiva, consider applying petroleum jelly with a cotton applicator for safety.
7. Dry affected tooth surfaces with triple syringe or if not feasible dry with cotton.
8. Bend microsponge, immerse into SDF, remove excess on side of dappen dish.
9. Apply directly onto the affected tooth surface(s) with microsponge.
10. Allow SDF to absorb for up to one minute if reasonable, then remove excess with gauze or cotton roll. (If using SSKI, apply with a different microsponge. Repeat one to three times until no further white precipitates are observed. Wait five to 10 seconds between applications. Remove excess with cotton.)
11. Rinse with water.
12. Place gloves, cotton and microbrushes into plastic waste bags.

FIGURE 3. Clinical protocol for the UCSF dental clinics.

over multiple years. We believe that dryness of the lesion during application is also important. Isolation with gauze and/or cotton rolls is sufficient, while air drying prior to application is thought to improve effectiveness. Allowing one to three minutes for the silver diamine fluoride to soak into and react with a lesion is thought to effect success.

Allowing only a few seconds to soak in due to the cooperation limits of very young patients commonly results in arrest. Application time in clinical studies does not correlate to outcome. However, our committee decided to be cautious in our recommendations for initial use. Longer absorption time also decreases concerns about removing silver

diamine fluoride with a posttreatment rinse. Removing any excess material with the same cotton used to isolate is routine to minimize systemic absorption.

Many clinicians place silver diamine fluoride at the diagnostic visit, then at one and/or three-month follow ups, then at semiannual recall visits (six, 12, 18, 24 months). Whether application needs

to continue after two or three years to maintain caries arrest is not known. Another approach is simply to substitute silver diamine fluoride for any application of fluoride varnish to a patient with untreated carious lesions. Increased frequency with higher disease burden follows the caries management by risk assessment (CAMBRA) principles.⁶¹ It is relevant to take photographs to track lesions over time.

Efforts to improve the penetration of silver diamine fluoride into affected dentin by chemical cavity preparation have not been studied but are being explored clinically. Pretreatment with ethylenediaminetetraacetic acid (EDTA) to remove superficial hydroxyapatite in affected dentin may open the dentinal tubules to further silver diamine fluoride penetration. Pretreatment with hypochlorite (bleach) may help breakdown bacteria and exposed dentin proteins, but this may be redundant to the action of the silver. Hypochlorite to decrease discoloration after silver diamine fluoride treatment is not recommended, as the color comes from silver that cannot be broken down like organic chromophores and might break down dentin proteins stabilized against the effects of bacteria and acid by interactions with silver.

Experience with the combination of silver nitrate plus fluoride varnish (see above) has many practitioners asking about a topical varnish after silver diamine fluoride placement to prevent silver diamine fluoride taste and keep the silver diamine fluoride in the lesion. We see no evidence that varnish would help achieve either goal. Varnish does not seal. Rather, allowing more time for residence and diffusion of silver diamine fluoride to react with and dry into the lesion is more likely to improve effectiveness. Also, in our experience, silver diamine fluoride results in less aversive taste and texture responses than to fluoride varnish.

Decreased darkening of lesions in the esthetic zone improves acceptance. SSKI is an option if the patient is not pregnant, though significant darkening should still be expected. SSKI and silver diamine fluoride are not to be combined prior to application — SSKI can be placed after drying the silver diamine fluoride-treated tooth. Silver diamine fluoride does not prevent restoration of a lesion, thus it does not prevent esthetic options. While silver diamine fluoride has been shown to be more effective than ART or interim restorative treatment

In our experience, silver diamine fluoride results in less aversive taste and texture responses than to fluoride varnish.

(IRT),³³ the two are compatible and can be combined across one or more visits.

The California Business and Professions Code permits dental hygienists and assistants to apply silver diamine fluoride for the control of caries because they are topical fluorides (Section 1910.(b)). Physicians, nurses and their assistants are permitted to apply fluorides in California and in many other states and federal programs. The recent decision of the Oregon Dental Board to allow dental hygienists and assistants to place silver diamine fluoride under existing rules for topical fluoride medicaments sets a precedent. Dental hygienists and assistants in Oregon were barred from providing silver nitrate in a previous decision. All providers need to be trained. Applications should be tracked if applied to the same patient by multiple clinics.

Documentation and Billing

A new code, D1354, for “interim caries arresting medication application” was approved by the Code on Dental Procedures and Nomenclature (CDT) Code Maintenance Commission for 2016. The code definition is “Conservative treatment of an active, nonsymptomatic carious lesion by topical application of a caries arresting or inhibiting medicament and without mechanical removal of sound tooth structure.” The CDT Code is the U.S. HIPAA standard code set and is required for billing. The Commission includes representatives from the major insurers, Medicaid, ADA, AGD and specialty organizations. Insurers are in the process of evaluating coverage for this treatment.

Legal Considerations

Silver diamine fluoride is cleared by the FDA for marketing as a Class II medical device to treat tooth sensitivity. We are discussing off-label use as a drug to treat and prevent dental caries. This is a parallel situation to fluoride varnish, which has the same device clearance but is ubiquitously used off label by dentists and physicians as a drug to prevent caries. The same public health dentists who achieved the FDA device clearance are now applying for a dental caries indication. However, this is a more complicated process, normally only carried out by large pharmaceutical companies, and is likely to take longer.

Consent

Because silver diamine fluoride is new in the U.S., it is important to communicate effectively. In the UCSF clinics, we are using a special consent form (**FIGURE 4**) as a way to inform patients, parents and caregivers, and

UCSF Dental Center Informed Consent for Silver Diamine Fluoride

Facts for consideration:

- Silver diamine fluoride (SDF) is an antibiotic liquid. We use SDF on cavities to help stop tooth decay. We also use it to treat tooth sensitivity. SDF application every six to 12 months is necessary.
- The procedure: 1. Dry the affected area. 2. Place a small amount of SDF on the affected area. 3. Allow SDF to dry for one minute. 4. Rinse.
- **Treatment with SDF does not eliminate the need for dental fillings or crowns to repair function or esthetics. Additional procedures will incur a separate fee.**
- I should not be treated with SDF if: 1. I am allergic to silver. 2. There are painful sores or raw areas on my gums (i.e., ulcerative gingivitis) or anywhere in my mouth (i.e., stomatitis).

Benefits of receiving SDF:

- SDF can help stop tooth decay.
- SDF can help relieve sensitivity.



Risks related to SDF include, but are not limited to:

- **The affected area will stain black permanently.** Healthy tooth structure will not stain. Stained tooth structure can be replaced with a filling or a crown.
- Tooth-colored fillings and crowns may discolor if SDF is applied to them. Color changes on the surface can normally be polished off. The edge between a tooth and filling may keep the color.
- If accidentally applied to the skin or gums, a brown or white stain may appear that causes no harm, cannot be washed off and will disappear in one to three weeks.
- You may notice a metallic taste. This will go away rapidly.
- If tooth decay is not arrested, the decay will progress. In that case the tooth will require further treatment, such as repeat SDF, a filling or crown, root canal treatment or extraction.
- These side effects may not include all of the possible situations reported by the manufacturer. If you notice other effects, please contact your dental provider.
- Every reasonable effort will be made to ensure the success of SDF treatment. There is a risk that the procedure will not stop the decay and no guarantee of success is granted or implied.

Alternatives to SDF, not limited to the following:

- No treatment, which may lead to continued deterioration of tooth structures and cosmetic appearance. Symptoms may increase in severity.
- Depending on the location and extent of the tooth decay, other treatment may include placement of fluoride varnish, a filling or crown, extraction or referral for advanced treatment modalities.

I CERTIFY THAT I HAVE READ AND FULLY UNDERSTAND THIS DOCUMENT AND ALL MY QUESTIONS WERE ANSWERED:

_____ (signature of patient) _____ (date)

_____ (signature of witness) _____ (date)

FIGURE 4. UCSF special consent form.

to standardize procedures because we have so many inexperienced student clinicians. All practices have established procedures for consent and an extra form may not be needed in the community. The normal elements of informed consent

apply. We sought to ensure awareness of the expected change in color of the dentin as the decay arrests, likelihood of reapplication and contraindications in the presence of silver allergy and stomatitis. Note the importance of distinguishing

between allergy to nickel and other trace metals rather than silver allergy, which is rare. We used readability measurements to guide intelligibility and included a progressively discoloring lesion to show stain of a lesion but not healthy enamel.

Conclusion

Silver diamine fluoride is a safe, effective treatment for dental caries across the age spectrum. At UCSF, it is indicated for patients with extreme caries risk, those who cannot tolerate conventional care, patients who must be stabilized so they can be restored over time, patients who are medically compromised or too frail to be treated conventionally and those in disparity populations with little access to care.

Application twice per year outperforms all minimally invasive options including the atraumatic restorative technique — with which it is compatible but 20 times less expensive. It approaches the success of dental fillings after two or more years, and again, prevents future caries — while fillings do not. Silver diamine fluoride is more effective as a primary preventive than any other available material, with the exception of dental sealants, which are > 10 times more expensive and need to be monitored.

Saliva may play a role in caries arrest by silver diamine fluoride. Lower rates of arrest are seen in geriatric patients.³⁸ The elderly tend to have less abundant and less functional saliva, which generally explains their higher caries rate. In pediatric patients, higher rates of arrest are noted for buccal or lingual smooth surfaces and anterior teeth.³¹ These surfaces bathe more directly in saliva than others. It is surprising that silver chloride is the main precipitant in treated dentin, as chloride is not a common component of dentin or silver diamine fluoride, so may come from the saliva.

Traditional approaches often provide only temporary benefit, given the highest rates of recurrent caries are in patients with the worst disease burden. The advent of a treatment for nonsymptomatic caries not requiring general anesthesia or sedation addresses long-standing

concerns about the expense, danger and practical complexity of these services.

Experience suggests that dryness prior to application enhances effectiveness. Good patient management is still profoundly relevant to the very young and otherwise challenged patients, though this one-minute intervention is more tolerable than other options. Silver diamine fluoride can readily replace fluoride varnish for the prevention of caries in patients who have active caries. This as a powerful new tool in the fight against dental caries, particularly suited for those who suffer most from this disease.

Clinical evidence supports continued application one to two times per year until the tooth is restored or exfoliates, and otherwise perhaps indefinitely. Some treated lesions keep growing, particularly those in the inner third of the dentin. It is unclear what will happen if treatment is stopped after two to three years and research is needed. ■

ACKNOWLEDGMENT

The UCSF paradigm shift committee subcommittee on silver caries arrest included Sean Mong, DDS, EdD; Spomenka Djordjevic, DDS, MS; Paul Atkinson, DDS, PhD; George Taylor, DMD, MPH, DrPh; Natalie Heavilin, DDS; Ling Zhan, DDS, PhD; John Featherstone, PhD; Hellen Ellenikiotis, DDS; and Jeremy Horst, DDS, PhD. Thanks to Linda Milgrom for designing the PubMed search. Thanks to Chad Zillich, DDS, for help with the literature review. Thanks to study authors, particularly Edward Lo, BDS, MDS, PhD; Chun Hung Chu, BDS, MDS, PhD; and Geoff Knight, BDS, MSc, MBA, for helpful discussions.

REFERENCES

- Clarkson BH, Exterkate RAM. Noninvasive dentistry: A dream or reality? *Caries Res* 2015;49 suppl 1(1):11-17. doi:10.1159/000380887.
- Yamaga R, Yokomizo I. Arrestment of caries of deciduous teeth with diamine silver fluoride. *Dent Outlook* 1969;33:1007-1013.
- Rosenblatt A, Stamford TCM, Niederman R. Silver diamine fluoride: A caries "silver-fluoride bullet." *J Dent Res* 2009;88(2):116-125. doi:10.1177/0022034508329406.
- Mei ML, Chu CH, Lo ECM, Samaranyake LP. Fluoride and silver concentrations of silver diamine fluoride solutions for dental use. *Int J Paediatr Dent* 2013;23(4):279-285. doi:10.1111/ipd.12005.
- Mei ML, Ito L, Cao Y, Li QL, Lo ECM, Chu CH. Inhibitory effect of silver diamine fluoride on dentine demineralisation and collagen degradation. *J Dent* 2013;41(9):809-817.

- doi:10.1016/j.jdent.2013.06.009.
- Hill TJ, Arnold FA. The effect of silver nitrate in the prevention of dental caries. I. The effect of silver nitrate upon the decalcification of enamel. *J Dent Res* 1937;16:23-28.
- Castillo JL, Rivera S, Aparicio T, et al. The short-term effects of diamine silver fluoride on tooth sensitivity: A randomized controlled trial. *J Dent Res* 2011;90(2):203-208. doi:10.1177/0022034510388516.
- Craig GG, Knight GM, McIntyre JM. Clinical evaluation of diamine silver fluoride/potassium iodide as a dentine desensitizing agent. A pilot study. *Aust Dent J* 2012;57(3):308-311. doi:10.1111/j.1834-7819.2012.01700.x.
- Markowitz K, Pashley DH. Discovering new treatments for sensitive teeth: The long path from biology to therapy. *J Oral Rehabil* 2008;35(4):300-315. doi:10.1111/j.1365-2842.2007.01798.x.
- Featherstone JDB. The continuum of dental caries — evidence for a dynamic disease process. *J Dent Res* 2004;83 Spec No C:C39-C42.
- Mei ML, Li QL, Chu CH, Yiu CKY, Lo ECM. The inhibitory effects of silver diamine fluoride at different concentrations on matrix metalloproteinases. *Dent Mater* 2012;28(8):903-908. doi:10.1016/j.dental.2012.04.011.
- Mei ML, Ito L, Cao Y, Li QL, Chu CH, Lo ECM. The inhibitory effects of silver diamine fluorides on cysteine cathepsins. *J Dent* 2014;42(3):329-335. doi:10.1016/j.jdent.2013.11.018.
- Klasen HJ. A Historical Review of the Use of Silver in the Treatment of Burns. II. Renewed Interest for Silver. 2000:131-138.
- Youravong N, Carlen A, Teanpaisan R, Dahlén G. Metal-ion susceptibility of oral bacterial species. *Left Appl Microbiol* 2011;53(3):324-328. doi:10.1111/j.1472-765X.2011.03110.x.
- Hamama HH, Yiu CK, Burrow MF. Effect of silver diamine fluoride and potassium iodide on residual bacteria in dentinal tubules. *Aust Dent J* 2015;60(1):80-87. doi:10.1111/adj.12276.
- Suzuki T, Nishida M, Sobue S, Moriwaki Y. Effects of diamine silver fluoride on tooth enamel. *J Osaka Univ Dent Sch* 1974;14:61-72.
- Chu CH, Lo ECM. Micro-hardness of dentin in primary teeth after topical fluoride applications. *J Dent* 2008;36(6):387-391. doi:10.1016/j.jdent.2008.02.013.
- Englander HR, James VE, Massler M. Histologic effects of silver nitrate of human dentin and pulp. *J Am Dent Assoc* 1958;57(5):621-630.
- Mei ML, Ito L, Cao Y, Lo ECM, Li QL, Chu CH. An *in vivo* study of arrested primary teeth caries with silver diamine fluoride therapy. *J Dent* 2014;42(4):395-402. doi:10.1016/j.jdent.2013.12.007.
- Knight GM, McIntyre JM, Craig GG, Mulyani, Zilm PS, Gully NJ. Inability to form a biofilm of *Streptococcus mutans* on silver fluoride- and potassium iodide-treated demineralized dentin. *Quintessence Int* 2009;40(2):155-161.
- Knight GM, McIntyre JM, Craig GG, Mulyani, Zilm PS, Gully NJ. An *in vitro* model to measure the effect of a silver fluoride and potassium iodide treatment on the permeability of demineralized dentine to *Streptococcus mutans*. *Aust Dent J* 2005;50(4):242-245.
- Knight GM, McIntyre JM, Craig GG, Mulyani, Zilm PS, Gully NJ. Differences between normal and demineralized

- dentine pretreated with silver fluoride and potassium iodide after an in vitro challenge by *Streptococcus mutans*. *Aust Dent J* 2007;52(1):16-21.
23. Wakshlak RB-K, Pedahzur R, Avnir D. Antibacterial activity of silver-killed bacteria: the "zombies" effect. *Sci Rep* 2015;5:9555. doi:10.1038/srep09555.
24. Duffin S. Back to the future: The medical management of caries introduction. *J Calif Dent Assoc* 2012;40(11):852-858.
25. Rethman MP, Beltrán-Aguilar ED, Billings RJ, et al. Nonfluoride caries-preventive agents: Executive summary of evidence-based clinical recommendations. *J Am Dent Assoc* 2011;142(9):1065-1071.
26. Plemons JM, Al-Hashimi I, Marek CL. Managing xerostomia and salivary gland hypofunction. *J Am Dent Assoc* 2014;145(8):867-873. doi:10.14219/jada.2014.44.
27. Stebbins EA. What value has argenti nitras as a therapeutic agent in dentistry? *Int Dent J* 1891;12:661-670.
28. Howe PR. A method of sterilizing and at the same time impregnating with a metal affected dentinal tissue. *Dent Cosmos* 1917;59:891-904.
29. Chu CH, Lo ECM, Lin HC. Effectiveness of silver diamine fluoride and sodium fluoride varnish in arresting dentin caries in Chinese preschool children. *J Dent Res* 2002;81(11):767-770.
30. Shah S, Bhaskar V, Venkataraghavan K, Choudhary P, Ganesh M, Trivedi K. Efficacy of silver diamine fluoride as an antibacterial as well as antiplaque agent compared to fluoride varnish and acidulated phosphate fluoride gel: An in vivo study. *Indian J Dent Res* 2013;24(5):575-581. doi:10.4103/0970-9290.123374.
31. Zhi QH, Lo ECM, Lin HC. Randomized clinical trial on effectiveness of silver diamine fluoride and glass ionomer in arresting dentine caries in preschool children. *J Dent* 2012;40(11):962-967. doi:10.1016/j.jdent.2012.08.002.
32. Yee R, Holmgren C, Mulder J, Lama D, Walker D, van Palenstein Helderma W. Efficacy of silver diamine fluoride for arresting caries treatment. *J Dent Res* 2009;88(7):644-647. doi:10.1177/0022034509338671.
33. Santos Dos VE, de Vasconcelos FMN, Ribeiro AG, Rosenblatt A. Paradigm shift in the effective treatment of caries in schoolchildren at risk. *Int Dent J* 2012;62(1):47-51. doi:10.1111/j.1875-595X.2011.00088.x.
34. Lo EC, Chu CH, Lin HC. A community-based caries control program for preschool children using topical fluorides: 18-month results. *J Dent Res* 2001;80(12):2071-2074.
35. Llodra JC, Rodriguez A, Ferrer B, Menardia V, Ramos T, Morato M. Efficacy of silver diamine fluoride for caries reduction in primary teeth and first permanent molars of schoolchildren: 36-month clinical trial. *J Dent Res* 2005;84(8):721-724.
36. Green E. A clinical evaluation of two methods of caries prevention in newly erupted first permanent molars. *Aust Dent J* 1989;34(5):407-409.
37. Craig GG, Powell KR, Price CA. Clinical evaluation of a modified silver fluoride application technique designed to facilitate lesion assessment in outreach programs. *BMC Oral Health* 2013;13(1):73. doi:10.1186/1472-6831-13-73.
38. Zhang W, McGrath C, Lo ECM, Li JY. Silver diamine fluoride and education to prevent and arrest root caries among community-dwelling elders. *Caries Res* 2013;47(4):284-290. doi:10.1159/000346620.
39. Liu BY, Lo ECM, Chu CH, Lin HC. Randomized trial on fluorides and sealants for fissure caries prevention *J Dent Res* 2012;91(8):753-758. doi:10.1177/0022034512452278.
40. Tan HP, Lo ECM, Dyson JE, Luo Y, Corbet EF. A randomized trial on root caries prevention in elders. *J Dent Res* 2010;89(10):1086-1090. doi:10.1177/0022034510375825.
41. Monse B, Heinrich-Weltzien R, Mulder J, Holmgren C, van Palenstein Helderma WH. Caries preventive efficacy of silver diamine fluoride (SDF) and ART sealants in a school-based daily fluoride toothbrushing program in the Philippines. *BMC Oral Health* 2012;12(1):52. doi:10.1186/1472-6831-12-52.
42. Li R, Lo ECM, Chu CH, Liu BY. Preventing and arresting root caries through silver diamine fluoride applications. In: 2014. iadr.confex.com/iadr/14iags/webprogram/Paper189080.html.
43. Duangthip D, Lo ECM, Chu CH. Arrest of dentin caries in preschool children by topical fluorides. In: 2014. iadr.confex.com/iadr/14iags/webprogram/Paper189031.html.
44. Frencken JE, Pilot T, Songpaisan Y, Phantumvanit P. Atraumatic Restorative Treatment (ART): Rationale, Technique and Development. *J Public Health Dent* 1996;56(3):135-140. doi:10.1111/j.1752-7325.1996.tb02423.x.
45. Innes NP, Evans DJ, Stirrups DR. The Hall Technique: A randomized controlled clinical trial of a novel method of managing carious primary molars in general dental practice: Acceptability of the technique and outcomes at 23 months. *BMC Oral Health* 2007;7(1):18. doi:10.1186/1472-6831-7-18.
46. Vasquez E, Zegarra G, Chirinos E, et al. Short-term serum pharmacokinetics of diamine silver fluoride after oral application. *BMC Oral Health* 2012;12(1):60. doi:10.1186/1472-6831-12-60.
47. Chu CH, Lo ECM. Promoting caries arrest in children with silver diamine fluoride: A review. *Oral Health Prev Dent* 2008;6(4):315-321.
48. Liu Y, Zhai W, Du F. [Clinical observation of treatment of tooth hypersensitivity with silver ammonia fluoride and potassium nitrate solution]. *Zhonghua Kou Qiang Yi Xue Za Zhi*. 1995;30(6):352-354.
49. Yamaga R, Nishino M, Yoshida S, Yokomizo I. Diamine silver fluoride and its clinical application. *J Osaka Univ Dent Sch* 1972;12(1):1-20.
50. Milgrom P, Taves DM, Kim AS, Watson GE, Horst JA. Pharmacokinetics of fluoride in toddlers after application of 5% sodium fluoride dental varnish. *Pediatrics* 2014;134(3):e870-e874. doi:10.1542/peds.2013-3501.
51. Goldberg ME, Cantillo J, Larijani GE, Torjman M, Vekeman D, Schieren H. Sevoflurane versus isoflurane for maintenance of anesthesia: Are serum inorganic fluoride ion concentrations of concern? *Anesth Analg* 1996;82(6):1268-1272.
52. Gotjamanos T, Afonso F. Unacceptably high levels of fluoride in commercial preparations of silver fluoride. *Aust Dent J* 1997;42(1):52-53.
53. Gotjamanos T, Ma P. Potential of 4 percent silver fluoride to induce fluorosis in rats: Clinical implications. *Aust Dent J* 2000;45(3):187-192.
54. Neesham DC. Fluoride concentration in AgF and dental fluorosis. *Aust Dent J* 1997;42(4):268-269.
55. Quock RL, Barros JA, Yang SW, Patel SA. Effect of Silver Diamine Fluoride on Microtensile Bond Strength to Dentin. *Oper Dent* 2012;37(6):610-616. doi:10.2341/11-344-L.
56. Knight GM, McIntyre JM, Mulyani. The effect of silver fluoride and potassium iodide on the bond strength of auto cure glass ionomer cement to dentine. *Aust Dent J* 2006;51(1):42-45.
57. Yamaga M, Koide T, Hieda T. Adhesiveness of glass ionomer cement containing tannin-fluoride preparation (HY agent) to dentin – an evaluation of adding various ratios of HY agent and combination with application diamine silver fluoride. *Dent Mater J* 1993;12(1):36-44.
58. Soeno K, Taira Y, Matsumura H, Atsuta M. Effect of desensitizers on bond strength of adhesive luting agents to dentin. *J Oral Rehabil* 2001;28(12):1122-1128.
59. Centers for Disease Control and Prevention. Long-Term Care Services in the United States: 2013 Overview, table 4. www.cdc.gov/nchs/data/nsllcp/long_term_care_services_2013.pdf.
60. U.S. Department of Health and Human Services. Integration of Oral Health and Primary Care Practice. www.hrsa.gov/publichealth/clinical/oralhealth. Accessed June 5, 2015.
61. Featherstone JDB, Adair SM, Anderson MH, et al. Caries management by risk assessment: Consensus statement, April 2002. *J Calif Dent Assoc* 2003 Mar;31(3):257-69.

THE CORRESPONDING AUTHOR, Jeremy A. Horst, DDS, PhD, can be reached at jeremy.horst@ucsf.edu.



CLINICAL USE OF SILVER DIAMINE FLUORIDE IN DENTAL TREATMENT

FEBRUARY 2016

Compendium of Continuing Education article provides an overview on the use of Silver Diamine Fluoride in dental treatment.

May L. Mei, BDS, MDS, PhD; Edward Chin-Man Lo, BDS, MDS, PhD; and Chun-Hung Chu, BDS, MDS, PhD.

ABSTRACT

The use of a topical fluoride solution, namely silver diamine fluoride (SDF), in dental treatment has been drawing increasing attention. SDF has been used in some countries in Asia, including Japan and China, as a caries-arresting and anti-hypersensitivity agent. It was recently cleared by the Food and Drug Administration in the United States as a fluoride to manage hypersensitive teeth. Topical application of SDF is a noninvasive procedure that is quick and simple to use. Promising results of laboratory studies and clinical trials have suggested that SDF is more effective than other fluoride agents to halt the caries process. A review concluded that SDF is a safe, effective, efficient, and equitable caries control agent that has a potentially broad application in dentistry and may meet the criteria of both the WHO Millennium Development Goals and the US Institute of Medicine's criteria for 21st century medical care. This article provides an overview of the clinical use of SDF in dental treatment.

Please use the link below to access the full article.

<https://cced.cdeworld.com/courses/4990#sthash.d0aJoy9Y.dpuf>



FRESH APPROACH TO CARIES ARREST IN ADULTS

SEPTEMBER/OCTOBER 2015

Decisions in Dentistry article on the use of Silver Diamine Fluoride in adult patients.

Dr. John Featherstone, Dean of the University of California San Francisco School of Dentistry and
Dr Jeremy Horst, DDS, PhD.

KEY TAKEAWAYS

- Cleared by the U.S. Food and Drug Administration for treating dentinal hypersensitivity, in off-label use silver diamine fluoride can be used to prevent and arrest caries.
- The agent acts as an antimicrobial that remains active well after application. It also promotes remineralization and resistance to demineralization in enamel and dentin.
- In order to effectively implement treatment, clinicians should know the indications and contraindications, and gain informed consent for use.
- Dentists and (if allowed by state practice acts) dental auxiliaries who apply this agent must understand precautions for handling silver diamine fluoride.
- Repeat application completely stops many, but not all lesions. Research is needed to determine why some caries are not arrested.

Please use the link below to access the full article.

<http://decisionsindentistry.com/article/fresh-approach-to-carries-arrest-in-adults/>

Additional Articles of Interest

1. Aust Dent J. 2015 Mar;60Suppl 1:95-105. doi: 10.1111/adj.12288. Improving the oral health of frail and functionally dependent elderly. Lewis A(1), Wallace J, Deutsch A, King P.
2. Int J Antimicrob Agents. 2015 Feb;45(2):183-7. doi: 10.1016/j.ijantimicag.2014.09.007. Epub 2014 Oct 14. Action of silver nanoparticles towards biological systems: cytotoxicity evaluation using hen's egg test and inhibition of *Streptococcus mutans* biofilm formation. FreirePL(1), Stamford TC(2), Albuquerque AJ(3), Sampaio FC(3), Cavalcante HM(3), Macedo RO(3), Galembeck A(4), Flores MA(5), Rosenblatt A(6).
3. Trials. 2014 Nov 19;15:448. doi: 10.1186/1745-6215-15-448. New proposal of silver diamine fluoride use in arresting approximal caries: study protocol for a randomized controlled trial. Mattos-Silveira J, Floriano I, Ferreira FR, Viganó ME, Frizzo MA, Reyes A, Novaes TF, Moriyama CM, Raggio DP, Imperato JC, Mendes FM, Braga MM(1).
4. Int J Pediatric Dent. 2014 Sep 17. doi: 10.1111/ipd.12134. [Epub ahead of print] Children's discomfort may vary among different treatments for initial approximal caries lesions: preliminary findings of a randomized controlled clinical trial. Mattos-SilveiraJ(1), Floriano I, Ferreira FR, Viganó ME, Mendes FM, Braga MM. Author information:
5. J Dent. 2014 Aug;42(8):945-51. doi: 10.1016/j.jdent.2014.05.017. Epub 2014 Jun 12. A new "silver-bullet" to treat caries in children--nano silver fluoride: a randomized clinical trial. Santos VE Jr(1), VasconcelosFilho A(2), Targino AG(3), Flores MA(4), Galembeck A(5), Caldas AF Jr(6), Rosenblatt A(7).
6. J Mater Sci Mater Med. 2014 Aug;25(8):2041-7. doi: 10.1007/s10856-014-5221-5. Epub 2014 May 13. An innovative approach to treating dental decay in children. A new anti-caries agent. TarginoAG(1), Flores MA, dos Santos Junior VE, de Godoy BenéBezerra F, de Luna Freire H, Galembeck A, Rosenblatt A.
7. Lasers Med Sci. 2015 Apr;30(3):985-91. doi: 10.1007/s10103-014-1521-8. Epub 2014 Jan 26. Effect of laser irradiation on the fluoride uptake of silver diamine fluoride treated dentine. Mei ML(1), Ito L, Zhang CF, Lo EC, Chu CH.
8. BMC Res Notes. 2014 Jan 3;7:3. doi: 10.1186/1756-0500-7-3. Arresting rampant dental caries with silver diamine fluoride in a young teenager suffering from chronic oral graft versus host disease post-bone marrow transplantation: a case report. Chu CH(1), Lee AH, Zheng L, Mei ML, Chan GC.
9. Org Lett. 2014 Jan 3;16(1):102-5. doi: 10.1021/ol403083e. Epub 2013 Dec 2. AgF-mediated fluorinativehomocoupling of gem-difluoroalkenes. GaoB(1),

Zhao Y, Ni C, Hu J.

10. Science. 2013 Nov 22;342(6161):956-60. doi: 10.1126/science. 1243759. Selective C-H fluorination of pyridines and diazines inspired by a classic amination reaction. FierPS(1), Hartwig JF.
11. Nat Protoc. 2013 Dec;8(12):2348-54. doi: 10.1038/nprot.2013.144. Epub 2013 Oct 31. Oxidative aliphatic C-H fluorination with manganese catalysts and fluoride ion. Liu W(1), Huang X, Groves JT.
12. J Dent. 2013 Sep;41(9):809-17. doi: 10.1016/j.jdent.2013.06.009. Epub 2013 Jun 27. Inhibitory effect of silver diamine fluoride on dentine demineralisation and collagen degradation. Mei ML(1), Ito L, Cao Y, Li QL, Lo EC, Chu CH.
13. Lasers Med Sci. 2014 Nov;29(6):1785-91. doi: 10.1007/s10103-013-1329-y. Epub 2013 Apr 26. Prevention of dentine caries using silver diamine fluoride application followed by Er:YAG laser irradiation: an in vitro study. Mei ML(1), Ito L, Chu CH, Lo EC, Zhang CF.
14. Spec Care Dentist. 2013 May-Jun;33(3):133-40. doi: 10.1111/j.1754-4505.2012.00318.x. Epub 2012 Dec 10. Prevention of root caries: a literature review of primary and secondary preventive agents. GluzmanR(1), Katz RV, Frey BJ, McGowan R.
15. Ann Clin Microbiol Antimicrob. 2013 Feb 26;12:4. doi: 10.1186/1476-0711-12-4. Antibacterial effects of silver diamine fluoride on multi-species cariogenic biofilm on caries. Mei ML(1), Li QL, Chu CH, Lo EC, Samaranayake LP.
16. Aust Dent J. 2013 Mar;58(1):50-6. doi: 10.1111/adj.12033. Epub 2013 Jan 30. An in vitro study of silver and fluoride ions on remineralization of demineralized enamel and dentine. ZhiQH(1), Lo EC, Kwok AC.
17. Contemp Clin Dent. 2012 Jul;3(3):262-4. doi: 10.4103/0976-237X.103615. Anti-microbial efficiency of silver diamine fluoride as an endodontic medication - An ex vivo study. Mathew VB(1), Madhusudhana K, Sivakumar N, Venugopal T, Reddy RK.
18. Am J Dent. 2012 Oct;25(5):299-302. Changes in the crystallinity of hydroxyapatite powder and structure of enamel treated with several concentrations of ammonium hexafluorosilicate. SugeT(1), Ishikawa K, Matsuo T.
19. J Oral Sci. 2012 Sep;54(3):267-72. Effects of ammonium hexafluorosilicate application on demineralized enamel and dentin of primary teeth. Ho-soyaY(1), Watanabe E, Tadokoro K, Inoue T, Miyazaki M, Tay FR.
20. Int J Paediatr Dent. 2013 Jul;23(4):279-85. doi: 10.1111/ipd.12005. Epub

- 2012 Oct 3. Fluoride and silver concentrations of silver diammine fluoride solutions for dental use. Mei ML(1), Chu CH, Lo EC, Samaranayake LP.
21. *In Vivo*. 2012 Jul-Aug;26(4):657-64. Effect of three fluoride compounds on the growth of oral normal and tumor cells. AcraAM(1), Sakagami H, Matsuta T, Adachi K, Otsuki S, Nakajima H, Koh T, Machino M, Ogiwara T, Watanabe K, Watanabe S, Salgado AV, Bastida NM.
 22. *J Dent Res*. 2012 Aug;91(8):753-8. doi: 10.1177/0022034512452278. Epub 2012 Jun 26. Randomized trial on fluorides and sealants for fissure caries prevention. Liu BY(1), Lo EC, Chu CH, Lin HC.
 23. *J Evid Based Dent Pract*. 2012 Jun;12(2):95-6. doi: 10.1016/j.jeb-dp.2012.03.011. Chlorhexidine varnish, sodium fluoride varnish, and silver diamine fluoride solution can prevent the development of new root caries in elders living in senior homes in Hong Kong. NiessenLC(1).
 24. *J Am Chem Soc*. 2012 Jul 4;134(26):10795-8. doi: 10.1021/ja304410x. Epub 2012 Jun 22. Copper-mediated fluorination of aryl iodides. Fier PS(1), Hartwig JF.
 25. *Oper Dent*. 2012 Nov-Dec;37(6):610-6. doi: 10.2341/11-344-L. Epub 2012 May 22. Effect of silver diamine fluoride on microtensile bond strength to dentin. Quock RL(1), Barros JA, Yang SW, Patel SA.
 26. *Am J Dent*. 2012 Feb;25(1):31-4. Antibacterial activity of ammonium hexa-fluorosilicate solution with antimicrobial agents for the prevention of dentin caries. Shibata S(1), Suge T, Kimura T, Ishikawa K, Matsuo T.
 27. *J Dent*. 2012 Jul;40(7):531-41. doi: 10.1016/j.jdent.2012.03.009. Epub 2012 Apr 3. Silver compounds used in dentistry for caries management: a review. PengJJ(1), Botelho MG, Matinlinna JP.
 28. *J Nanosci Nanotechnol*. 2011 Nov;11(11):10063-8. Fabrication and enhanced visible light photocatalytic activity of fluorine doped TiO₂ by loaded with Ag. Lin X(1), Rong F, Ji X, Fu D, Yuan C.
 29. *Aust Dent J*. 2012 Mar;57(1):65-70. doi: 10.1111/j. 1834-7819.2011.01641.x. Effect of silver and fluoride ions on enamel demineralization: a quantitative study using micro-computed tomography. Liu BY(1), Lo EC, Li CM.
 30. *Acta Odontol Latinoam*. 2011;24(2):127-31. In vitro antibacterial activity of silver diamine fluoride in different concentrations. de Almeida LdeF(1), Cavalcanti YW, Valença AM.
 31. *J Calif Dent Assoc*. 2011 Oct;39(10):735-41. Prevention-centered caries management strategies during critical periods in early childhood. MilgromP(1), Chi DL.

32. J Conserv Dent. 2011 Jul;14(3):233-6. doi: 10.4103/0972-0707.85796. Remineralizing efficacy of silver diamine fluoride and glass ionomer type VII for their proposed use as indirect pulp capping materials - Part II (A clinical study). SinhaN(1), Gupta A, Logani A, Shah N.
33. J Org Chem. 2011 Oct 21;76(20):8543-8. doi: 10.1021/jo2016168. Epub 2011 Sep 26. Palladium-catalyzed direct ortho C-H arylation of 2-arylpyridine derivatives with aryltrimethoxysilane. Li W(1), Yin Z, Jiang X, Sun P.
34. J Conserv Dent. 2011 Apr;14(2):113-6. doi: 10.4103/0972-0707.82603. An ex vivo study to evaluate the remineralizing and antimicrobial efficacy of silver diamine fluoride and glass ionomer cement type VII for their proposed use as indirect pulp capping materials - Part I. Gupta A(1), Sinha N, Logani A, Shah N.
35. J Dent. 2011 Sep;39(9):612-8. doi: 10.1016/j.jdent.2011.06.008. Epub 2011 Jul 1. Reaction of silver diamine [corrected] fluoride with hydroxyapatite and protein. Lou YL(1), Botelho MG, Darvell BW.
36. Int J Paediatr Dent. 2012 Jan;22(1):2-10. doi: 10.1111/j.1365-263X.2011.01149.x. Epub 2011 Jun 27. Effects of silver diamine fluoride on dentine carious lesions induced by Streptococcus mutans and Actinomyces naeslundii biofilms. Chu CH(1), Mei L, Seneviratne CJ, Lo EC.
37. J Org Chem. 2011 Jul 15;76(14):5793-802. doi: 10.1021/jo200966k. Epub 2011 Jun 21. Synthesis of triazafluoranthrenones via silver(I)-mediated non-oxidative and oxidative intramolecular palladium-catalyzed cyclizations. Koutentis PA(1), Loizou G, Lo Re D.
38. Med Hypotheses. 2011 Sep;77(3):315-7. doi: 10.1016/j.mehy.2011.05.002. Epub 2011 May 19. Is a drill-less dental filling possible? QuockRL(1), Patel SA, Falcao FA, Barros JA.
39. J Evid Based Dent Pract. 2011 Mar;11(1):54-5. doi: 10.1016/j.jeb-dp.2010.11.011. Topical application of silver diamine fluoride may arrest dental caries. HouptM(1).
40. Org Lett. 2011 Mar 18;13(6):1510-3. doi: 10.1021/ol200196m. Epub 2011 Feb 11. Silver-mediated palladium-catalyzed direct C-H arylation of 3-bromoisothiazole-4-carbonitrile. IoannidouHA(1), Koutentis PA.
41. J Mol Model. 2011 Sep;17(9):2237-48. doi: 10.1007/s00894-010-0949-4. Epub 2011 Jan 22. On possible existence of pseudobinary mixed valence fluorides of Ag(I)/Ag(II): a DFT study. Grochala W(1).
42. Anal Chem. 2011 Feb 15;83(4):1315-20. doi: 10.1021/ac1024683. Epub 2011 Jan 14. Quantification of VX vapor in ambient air by liquid chromatography isotope dilution tandem mass spectrometric analysis of glass bead filled sam-

- pling tubes. Evans RA(1), Smith WL, Nguyen NP, Crouse KL, Crouse CL, Norman SD, Jakubowski EM.
43. Stat Med. 2011 Feb 10;30(3):250-9. doi: 10.1002/sim.4094. Epub 2010 Nov 5. Analysis of multilevel grouped survival data with time-varying regression coefficients. Wong MC(1), Lam KF, Lo EC.
 44. MolPharmacol. 2010 Nov;78(5):952-60. doi: 10.1124/mol.110.066407. Epub 2010 Aug 20. Inhibition of large-conductance Ca²⁺-activated K⁺ channels by nanomolar concentrations of Ag⁺. Zhou Y(1), Xia X, Lingle CJ.
 45. Zhongguo Yi XueKeXue Yuan XueBao. 2010 Jun;32(3):265-8. doi: 10.3881/j.issn.1000-503X.2010.03.005. [Effect of combination of pulsed CO₂ laser irradiation and diammine silver fluoride treatment on ultrastructure of dentine]. [Article in Chinese] Wang Q(1), Zhao JZ, Wu XM.
 46. J Endod. 2010 Jun;36(6):1026-9. doi: 10.1016/j.joen.2010.02.029. Epub 2010 Apr 24. Antimicrobial efficacy of 3.8% silver diamine fluoride and its effect on root dentin. HiraishiN(1), Yiu CK, King NM, Tagami J, Tay FR.
 47. J Evid Based Dent Pract. 2010 Jun;10(2):122-4. doi: 10.1016/j.jebdp.2010.02.014. Silver diamine fluoride (SDF) may be better than fluoride varnish and no treatment in arresting and preventing cavitated carious lesions. Beltrán-Aguilar ED(1).
 48. Dent Today. 2010 Feb;29(2):130, 132-3. Leave decay in my cavity? You must be kidding! Knight GM(1), McIntyre JM, Craig GG, Mulyani.
 49. J Am Chem Soc. 2010 Feb 10;132(5):1476-7. doi: 10.1021/ja909806t. Silver dendrites from galvanic displacement on commercial aluminum foil as an effective SERS substrate. GutiérrezA(1), Carraro C, Maboudian R.
 50. Braz Oral Res. 2009 Jul-Sep;23(3):296-301. In vitro evaluation of fluoride products in the development of carious lesions in deciduous teeth. SantosLdeM(1), Reis JI, Medeiros MP, Ramos SM, Araújo JM.
 51. AcadPediatr. 2009 Nov-Dec;9(6):404-9. doi: 10.1016/j.acap.2009.09.001. An examination of the advances in science and technology of prevention of tooth decay in young children since the Surgeon General's Report on Oral Health. MilgromP(1), Zero DT, Tanzer JM.
 52. Evid Based Dent. 2009;10(3):68. doi: 10.1038/sj.ebd.6400661. Silver lining for caries cloud? DeeryC(1).
 53. J Org Chem. 2009 Nov 6;74(21):8232-42. doi: 10.1021/jo901725k. 5(6)-anti-Substituted-2-azabicyclo[2.1.1]hexanes: a nucleophilic displacement route. KrowGR(1), Edupuganti R, Gandla D, Choudhary A, Lin G, Sonnet PE, DeBrosse C, Ross CW 3rd, Cannon KC, Raines RT.

54. Dent Mater. 2010 Jan;26(1):29-34. doi: 10.1016/j.dental. 2009.08.011. Effects of ammonium hexafluorosilicate concentration on dentin tubule occlusion and composition of the precipitate. SugeT(1), Kawasaki A, Ishikawa K, Matsuo T, Ebisu S.
55. J Dent Child (Chic). 2009 Jan-Apr;76(1):28-33. Effect of silver diammine fluoride on incipient caries lesions in erupting permanent first molars: a pilot study. Braga MM(1), Mendes FM, De Benedetto MS, Imperato JC.
56. Journal of Advanced Oral Research / Jan-Apr 2014 / Vol. 5 No.1. Silver Diamine Fluoride: A Review and Current Applications. Shalin Shah¹, Vijay Bhaskar², Karthik Venkatraghavan³, Prashant Choudhary⁴, Ganesh M.⁵, Krishna Trivedi⁶
57. International Dental Journal 2012; 62: 47–51. doi: 10.1111/j.1875-595X.2011.00088.x. Paradigm shift in the effective treatment of caries in schoolchildren at risk. Valdeci E. dos Santos Jr, Fla´via M. N. de Vasconcelos¹, Andre´a G. Ribeiro and Aronita Rosenblatt
58. J Dent. 2015 May 30. pii: S0300-5712(15)00124-4. doi: 10.1016/j.jdent.2015.05.006. A randomized clinical trial on arresting dentine caries in preschool children by topical fluorides -18 month results. Duangthip D, Chu CH, Lo CM
59. Parental Acceptance of the Use of Diamine Silver Fluoride in Children Aged 0 to 3 Years in the City of Cascavel, PR, Brazil Thaisa Cez´aria TRICHES, Mabel Mariela Rodr´ıguez CORDEIRO, Juliana Garcia Mugnai Vieira SOUZA, Eduardo Karam SALTORI, Beatriz Helena Sottile FRANÇA
60. Pulp response to high fluoride releasing glass ionomer, silver diamine fluoride, and calcium hydroxide used for indirect pulp treatment: An in-vivo comparative study. AtishKorwar, Sidhartha Sharma, Ajay Logani, and Naseem-ShahContempClin Dent. 2015 Jul-Sep; 6(3): 288–292.
61. The evidence base for professional and self-care prevention - caries, erosion and sensitivity. Twetman S. BMC Oral Health. 2015;15 Suppl 1:S4. doi: 10.1186/1472-6831-15-S1-S4. Epub 2015 Sep 15.
62. Root Caries in Older Adults. Gregory D, Hyde S. J Calif Dent Assoc. 2015 Aug;43(8):439-45.
63. Silver diamine fluoride and glass ionomer differentially remineralize early caries lesions, in situ. Nantanee R, Santiwong B, Trairatvorakul C, Hamba H, Tagami J. Clin Oral Investig. 2015 Sep 23.
64. The effectiveness of the biannual application of silver nitrate solution followed by sodium fluoride varnish in arresting early childhood caries in preschool children: study protocol for a randomised controlled trial. Chu

CH, Gao SS, Li SK, Wong MC, Lo EC. *Trials*. 2015 Sep 25;16(1):426. doi: 10.1186/s13063-015-0960-2

65. Prevention of secondary Caries by silver diamine fluoride. Mei ML, Zhao IS, Ito L, Lo EC, Chu CH. *Int Dent J*. 2015 Dec 22. doi: 10.1111/idj.12207
66. An alternate technique of care using silver fluoride followed by stannous fluoride in the management of root caries in aged care. Deutsch A. *Spec Care Dentist*. 2015 Dec 21. doi: 10.1111/scd.12153.
67. Caries remineralisation and arresting effect in children by professionally applied fluoride treatment - a systematic review. Gao SS, Zhang S, Mei ML Lo EC, Chu CH. *BMC Oral Health*. 2016 Feb 1; 16(1), 12.
68. Rosenblatt A, Stamford TCM, Niederman R. Silver diamine fluoride: a caries “silver-fluoride bullet.” *J Dent Res*. 2009;88(2):116–125.

Frequently Asked Questions

1. **Should SDF be used as a caries preventive therapy in a similar manner to how fluoride varnish is used “off-label” as a caries prevention therapy?**

No. SDF is indicated for site-specific application only. Fluoride varnish can be applied to specific sites, but most fluoride varnish applications are full mouth. SDF is never a full mouth application.

SDF treatment should be limited to up to five sites per appointment. Application to more than five sites should be spaced by 7 days. For patients with active caries, SDF can be considered as a replacement of fluoride varnish as the primary means to prevent new lesions, with application of SDF to the active lesions only.

2. **Is SDF safe for use in children?**

Regarding the margin of safety for dosing, a study was conducted for FDA review for market clearance in rats and mice to determine the lethal dose by oral and subcutaneous administration. The worst case scenario is subcutaneous administration and that lethal dose was found to be 380 mg/kg. One drop (25uL) of 38% silver diamine fluoride (SDF) contains 9.5 mg silver diamine fluoride. Thus, one drop of 38% SDF applied to 10 kg (22 lb.) child would equal 0.95 mg/kg, equal to a four-hundred fold safety margin.

In setting up protocols for undergraduate application of 38% SDF the University of California San Francisco set a recommended limit of one drop per 10 kg per treatment visit, with weekly intervals at most.

3. **Does SDF discolor skin or oral tissue?**

Contact to skin is not harmful but is likely to cause temporary “tattooing”. The effect is not immediate, rather it will be noticed within hours. The speed of discoloration is accelerated with light contact. The staining will be limited to direct areas of contact and will fade over a period of 24-72 hours. Patients should be protected with bibs and safety glasses as in any clinical procedure. If you believe you have touched the applicator to the skin of a patient it is good to advise them of possible tattooing.

Contact to oral soft tissue is less likely to cause tattooing, but is still possible. Take care to protect soft tissue with petroleum jelly or cocoa butter when applications is adjacent to gingival tissue (root caries, treatment of restoration margins). Light blanching is also possible from prolonged direct contact, but has been reported to be minor and resolves itself within 1-2 days.

4. Does the application technique differ between the label indication of relief of dentinal hypersensitivity and the off label indication of caries control?

In countries where SDF is used for the control of caries the application technique is identical to the instructions we provide in the Advantage Arrest Package Insert for the relief of dentinal hypersensitivity. No excavation, decay removal or anesthesia is required. The area to be treated should be “dry tooth brush” clean, free of plaque and debris. The area should be free of saliva, so as not to dilute the material. Transfer the material from a disposable plastic dappen dish to the surface to be treated and allow time to dry, which generally occurs in 30-60 seconds. If accelerated drying is required due to patient compliance use a low/weak air stream to dry the material.

The chemical action of the SDF occurs almost immediately in the outer layers of the softened dentin and can be confirmed by changes in the hardness and density of the dentin surface, similar to caries that arrests naturally because of positive changes in oral hygiene, diet, or daily application of fluoride in custom trays. The darkening of the lesion occurs over 24 hours and may increase over a week. Reexamination of the lesion at the next regular recall is appropriate and reapplication of SDF may be warranted. Repeat until the lesion has arrested

5. Is there a recommended frequency of application of SDF for caries control?

Caries arrest studies were conducted with SDF applications of once and twice annually. Arrested lesions were retreated every six-months.

Clinicians have reported that they will recall their first cohort of SDF patients within 3-6 weeks to evaluate the application and action of the treatment. Once they have a feel for the predictability of the material with their application technique they will set recall appointments based on the risk level and caries activity of the patient with higher risk patients at 3-month intervals. Moderate to high-risk patients, where it appears that home care and diet counseling has had positive impact, are recalled at 6-months.

6. Does the application of SDF to a lesion cause discoloration?

Yes, darkening of decayed, demineralized sites occurs as the lesion arrests. This is similar to what is seen when caries arrests from changes in diet or increased use of other fluorides. A recent study showed that patients see the discoloration as a clear indication that the treatment is working. Similar to the treatment of eroded and hypersensitive dentin, the treated area can be restored using glass ionomer or with a sandwich restoration of both glass ionomer and composite. In lab studies, bond strength of composite by itself to treated surfaces is reduced but the clinical importance is unknown.

38% silver diamine fluoride should not be diluted in an attempt to reduce discoloration. Studies have shown that diluted solutions may not be effective for caries arrest.

Ionic silver adsorbs onto almost any protein surface and is especially tenaciously bound to denatured proteins. This accounts for the specificity to carious collagen over normal collagen, but both will stain. The differentiator between these stains is that with SDF use, intrinsic pigmentation of a carious lesion occurs and surface protein staining occurs primarily on healthy tissue. These oxides are bound to the tissue and don't wash or polish away. This is why the blackened lesion retains its dark color for so long, and is most likely the reason the antimicrobial effect is long lasting.

The functional indicator of effectiveness is when the silver oxide is bound to the diseased collagen. If the surface doesn't turn black, the silver didn't bind and the antimicrobial effect will only be short lived.

7. Can SDF be used on a prepared tooth just prior to restoration cementation?

Desensitizing agents have been shown to be protective of the pulp when placed on crown preparations to reduce dentin permeability. Advantage Arrest, a desensitizer, has been shown safe to the pulp when placed on exposed dentin. In addition, studies have shown desensitization and efficacy in treating softened dentin before placing direct restorations. Usually the tooth is first treated with silver diamine fluoride 38%. This provides the benefit of sealing tubules plus the antimicrobial benefits of both silver and fluoride

8. Does an SDF treated site compromise the bond strength of glass ionomer (GI), resin-modified glass ionomer (RMGI) or resin composite restorations?

A recent in vitro study investigated the micro-tensile bonding strength of resin composite to the dentin of primary molars and found that pretreating does not affect the bonding strength. The study concluded: "In the SDF group, the fracture occurred most often within the adhesive layer, suggesting that bonding strength might be stronger between the adhesive and the dentin pretreated with SDF." (Pediatric Dentistry, V 38, N 2, Mar/Apr 2016, pgs. 148-153)

The recent January 2016 SDF review in the California Dental Association Journal stated that: "Using a contemporary bonding system, silver diamine fluoride had no effect on composite bonding to non-carious dentin using either self-etch or full-etch systems. In one study, simply rinsing after silver diamine fluoride application avoided a 50 percent decrease in bond strength for GIC. In another study, increased dentin bond strength to GIC was observed. Silver diamine fluoride decreased dentin bonding strength of resin-based crown cement by approximately one-third. Thus, rinsing will suffice for direct restorations, while excavation of the silver diamine fluoride-treated superficial dentin is appropriate for cementing crowns.

SDF treated sites tend to discolor more rapidly with light curing. Care should

be taken when bonding translucent restorative materials in anterior teeth. The use of opaquers is recommended when covering extensive anterior treated sites. Self cured materials may diminish anterior discoloration issues associated with light curing.

9. Are there any contraindications for the use of SDF for the control of caries?

SDF should not be placed on exposed pulps. Other topical fluorides (e.g. fluoride varnish) should not be used in the same appointment. Studies have shown that 38% silver diamine fluoride conveys more effective protection against decay in other teeth than fluoride varnish with reduced overall fluoride exposure.

10. Is there evidence of caries prevention benefit to non-application sites following SDF use for a patient?

Treating carious areas with silver diamine fluoride 38% acts as a whole mouth fluoride treatment. It can also be used in place of sealants in grooves. A protective effect has been shown to non-treated teeth and surfaces. These findings come from high quality randomized clinical trials.

11. Are there any post appointment instructions for the patient or the caregivers/guardians?

There are no postoperative limitations. Patients may eat or drink immediately. Patients may brush their teeth with fluoridated toothpaste on their regular schedule.

12. Does SDF stain countertops, instruments etc.?

Yes. When dispensing SDF it is a good idea to use an absorbent material that has a coated bottom, like a patient bib, under the dappen dish and applicator to avoid contact with metal trays and office countertops. If SDF comes in contact with instruments or countertops wash immediately with water, soap, ammonia or iodine tincture and then rinse thoroughly with water. Sodium hypochlorite (household bleach) can also be used for difficult stains.

13. What are the safety implications for application of SDF for a patient that has more than six sites to be treated?

The Margin of Safety for the volume of product needed to treat six sites is within 130 times the NOAEL (no-observed-adverse-effect-level). Treating more sites in one visit will likely have little practical impact on patient safety. Like protocols for fluoride varnish application, the suspension for several days of fluoride supplements is advised.

14. How does an arrested lesion treated with SDF look like on radiographs?

Arrested lesions look like a scar on radiographs. You will observe radio-opacity as the mineralization of the previously softened dentin increases. Ultimately the best test of arrest is still the color change and tactile hardness of the dentin surface.

It is advised that you educate your referring dentist about your use of Advantage Arrest since the appearance of a treated lesion might be new and confusing for many practitioners.

15. How can Advantage Arrest be coded using CDT?

There is a new CDT code for 2016 specifically for the use of caries arresting medicaments; the off-label use of Advantage Arrest.

Code D1354

The nomenclature reads: “Interim caries arresting medicament application,” with the descriptor; “Conservative treatment of an active, non-symptomatic carious lesion by topical application of a caries arresting or inhibiting medicament and without mechanical removal of sound tooth structure.”

It is common for insurance providers to initially not reimburse for new codes as they are developing usual and customary rates for the procedure. However, it’s important the new code is used so the providers can see the volume of use and determine future coverage. There are several providers that have announced coverage in various states.

For a current list of providers visit: https://en.wikipedia.org/wiki/Silver_diamine_fluoride

There are three other options to code the use of Advantage Arrest Silver Diamine Fluoride 38%. These codes are:

D1208 - Topical application of fluoride

SDF is categorized as a fluoride and can be used to treat site-specific locations under this code if there is no active lesion present. It’s application and effect is very different than most fluorides.

D 9910 - Application of a desensitizing medicament, per visit

SDF is indicated for dentinal hypersensitivity treatment and can be used to treat site-specific locations.

D1999 - Unspecified preventive procedure by report

From a third party payer perspective, this is the preferred code so providers can track the frequency of a procedure and develop usual and customary rates for future coverage.

It is also helpful to identify caries risk to justify the reimbursement with a recognized caries risk tool. Codes: D0601 (low), D0602 (moderate) and D0603 (high) codes are especially helpful in adult claims.

16. Can SDF be used as a cavity Liner?

SDF is cleared in the same FDA category as cavity liners. Although there are no head to head clinical trials comparing SDF as a cavity liner, it has been used successfully in this way.

SDF will not discolor intact enamel or dentin. SDF can discolor demineralized tooth structure brown/black. Some of this discoloration may shadow a restoration and can create less than optimal esthetic restorations.

17. Who is allowed to apply SDF in clinical practice in my State?

Each State dental practice act is different. Since SDF is a fluoride containing product indicated for the control of dentinal hypersensitivity it should fit into the same rules as fluoride varnishes. Please confirm that within your own State's dental practice acts.

18. How far into enamel and dentin does SDF penetrate?

Approximately 25 microns into enamel and 300 microns into dentin. This will seal off the surface of any lesions and cause the remainder of the lesion to arrest. In a 2002 study by Dr. Chu, 100% of lesions stained black to the outer edge of the lesion were arrested.

Breakthrough Therapy Designation for Silver Diamine Fluoride granted by FDA

- **Advantage Arrest™ Silver Diamine Fluoride 38% is the first drug to receive Breakthrough Therapy Designation for the arrest of tooth decay**
- **It is the only Breakthrough Therapy Designation ever received for an oral care medicine**

Redmond, OR and West Palm Beach, FL – October 30, 2016 -- Advantage Silver Dental Arrest, LLC and Elevate Oral Care, LLC announced today that the U.S. Food and Drug Administration (FDA) has granted “*Breakthrough Therapy Designation*” to *Advantage Arrest™ Silver Diamine Fluoride 38%* for the arrest of tooth decay in children and adults.

Breakthrough Therapy Designation represents the FDA’s effort to address an unmet, serious, life-threatening medical need where there is no available therapy. It is a process designed to expedite the development and review of drugs that are intended to treat a serious condition as soon as possible and preliminary clinical evidence indicates that the drug may demonstrate substantial improvement over available therapies on a clinically significant endpoint(s).

In use for over 80 years as a caries-arresting drug in Japan and other countries, the Designation was granted based on the FDA review of 10 worldwide randomized clinical trials evaluating silver diamine fluoride for caries arrest in children aged 3-9 or adults aged 60-89, as well as studies conducted by Advantage Silver Dental Arrest, LLC. Collectively these trials involved approximately 1500 subjects who were monitored for 1-3 years.

Currently, Advantage Arrest Silver Diamine Fluoride 38% is cleared by the FDA as a medical device for the treatment of dentinal hypersensitivity. It is well known that since it received FDA clearance in 2014 and became commercially available in April 2015, the material has been used extensively by oral health professionals in the U.S. off-label for the arrest of dental caries.

The American Dental Association added a new Code on Dental Procedures and Nomenclature (CDT Code D1354 – Interim Caries Arresting Medicament Application) to achieve uniformity, consistency and specificity in accurately documenting treatment for the efficient processing of dental claims. Multiple insurers and State Medicaid programs are currently reimbursing for the cost of the therapy for caries arrest.

“The Breakthrough Therapy Designation represents a major milestone in our over eight year effort, spearheaded by Dr. Peter Milgrom of Advantage and the University of

Washington, to improve the treatment of severe early childhood dental caries, a serious and rampant disease,” said Dr. Mike Shirtcliff, President of Advantage Silver Dental Arrest, LLC.

Kevin Thomas, President of Elevate Oral Care, the commercialization organization for Advantage Arrest Silver Diamine Fluoride 38% stated, “We are proud to be a part of this exciting advancement that fills such a huge unmet need.”

About Advantage Silver Dental Arrest, LLC

Advantage Silver Dental Arrest (ASDA) is a limited liability company headquartered in Redmond, Oregon. ASDA develops innovative drug products and medical devices for improving oral health. Its professional staff includes internationally recognized scientists with a long record of research achievements and contributions in oral care. In addition to receiving FDA Breakthrough Therapy Designation for silver diamine fluoride solution, ASDA has received multiple Small Business Innovation Research (SBIR) grants for its work in developing innovative dental care technologies.

About Elevate Oral Care, LLC

Founded six years ago, Elevate Oral Care develops, manufactures and markets oral preventive care products for healthcare professionals and consumers. The Elevate management team consists of the former principal owners of OMNII Oral Pharmaceuticals, which was acquired by 3M Company in 2005, as well as a number of other oral care industry professionals with a combined 100 years plus experience in the oral care industry. The current Elevate product line includes our patented FluoriMax® 2.5% Sodium Fluoride Varnish, patent-pending Allday™ Dry Mouth Spray and our proprietary Just Right® metered-dose children’s toothpaste.

Contacts:

Advantage Silver Dental Arrest, LLC
Cheryl Barker, 541-504-3945
cherylb@advantagedental.com

Elevate Oral Care, LLC
Kevin Thomas, 877-866-9113
kthomas@elevateoralcare.com

Sapp, Jessica

From: Tomar, Scott <STOMAR@dental.ufl.edu>
Sent: Wednesday, September 14, 2016 8:24 AM
To: Sapp, Jessica
Subject: Silver diamine fluoride application by dental hygienists

Dear Ms. Sapp,

I'm sorry we haven't been able to connect by telephone, but thank you for providing your email address.

First, some background:

As you probably are aware, silver diamine fluoride (SDF) was approved by the US Food & Drug Administration for use in treating dentinal hypersensitivity (the same indication for which fluoride varnish was approved). The product, Advantage Arrest, is manufactured by Elevate Oral Care and is now available throughout the country. Similar to fluoride varnish, SDF is being used off-label for management of dental caries and its use is being taught in dental schools. Numerous articles on the use of SDF for caries management have appeared in US dental journals.

My specific question:

In Florida, dental hygienists may perform their duties in health access settings, and they may "Apply topical fluorides, including fluoride varnishes, which are approved by the American Dental Association or the Food and Drug Administration." Is a dental hygienist in Florida working in a health access setting allowed to apply silver diamine fluoride?

Thank you,

Scott Tomar

Scott L. Tomar, DMD, MPH, DrPH
Professor & Interim Chair
Dept. of Community Dentistry & Behavioral Science, College of Dentistry
University of Florida
1329 SW 16th Street, Suite 5180
PO Box 103628
Gainesville, FL 32610-3628
Tel: 352.273.5968 / Fax: 352.273.5985
stomar@dental.ufl.edu

Florida Department of Education
Curriculum Framework

Program Title: Dental Hygiene
Career Cluster: Health Science

AS

CIP Number	1351060200
Program Type	College Credit
Standard Length	88 credit hours
CTSO	HOSA: Future Health Professionals
SOC Codes (all applicable)	29-2021 Dental Hygienists
CTE Program Resources	http://www.fldoe.org/academics/career-adult-edu/career-tech-edu/program-resources.stml

Purpose

This program offers a sequence of courses that provides coherent and rigorous content aligned with challenging academic standards and relevant technical knowledge and skills needed to prepare for further education and careers in the Health Science career cluster; provides technical skill proficiency, and includes competency-based applied learning that contributes to the academic knowledge, higher-order reasoning and problem-solving skills, work attitudes, general employability skills, technical skills, and occupation-specific skills, and knowledge of all aspects of Health Science career cluster.

This program is designed to prepare students for employment as dental hygienists SOC Code-29-2021 Dental Hygienist or to provide supplemental training for persons previously or currently employed in this occupation.

The content includes but is not limited to patient assessment, dental hygiene instrumentation and direct patient care services (scaling/root planing/curettage/radiographs/oral hygiene-instruction/expanded functions), community dental health, dental office emergencies, infection control, special needs dental care, office management, employability skills, ethics and jurisprudence.

Additional Information relevant to this Career and Technical Education (CTE) program is provided at the end of this document.

Program Structure

This program is a planned sequence of instruction consisting of 88 credit hours.

Regulated Programs

Students are prepared to take the Dental Hygiene National Board and state licensure examinations. Dental Hygiene Programs accredited by the American Dental Association Commission on Dental Accreditation are required to implement clinical experiences outlined in these program standards.

Standards

After successfully completing this program, the student will be able to perform the following:

- 01.0 Demonstrate knowledge of the dental health care delivery system and dental health occupations
- 02.0 Use oral and written communication skills in creating, expressing and interpreting information and ideas
- 03.0 Describe the legal and ethical responsibilities of the dental health care worker
- 04.0 Demonstrate an understanding of general anatomy and physiology and apply wellness and disease concepts
- 05.0 Demonstrate the importance of health, safety, and environmental management systems in dental organizations and their importance to organizational performance and regulatory compliance
- 06.0 Recognize and respond to emergency situations
- 07.0 Use information technology tools
- 08.0 Explain the importance of employability skills
- 09.0 Demonstrate knowledge of blood borne diseases, including HIV/AIDS
- 10.0 Demonstrate leadership and teamwork skills needed to accomplish team goals and objectives
- 11.0 Perform expanded functions for the dental hygienist as permitted by Florida Statutes/Law
- 12.0 Perform dental office procedures
- 13.0 Identify, describe, maintain and utilize dental instruments and equipment
- 14.0 Identify and perform dental and radiographic procedures
- 15.0 Identify properties and uses, and manipulate dental materials
- 16.0 Describe the legal and ethical responsibilities of the dental hygienist
- 17.0 Identify and explain the formation and function of the head, neck, dental structures and tissues including pathological conditions of the human body in relation to the oral cavity
- 18.0 Identify and explain principles of microbiology, disease transmission, disease prevention, and perform infection control procedures
- 19.0 Identify and explain usage, administration, indications, contraindications, adverse reactions and precautions of pharmaceutical and anesthetic agents used in the treatment of dental disease
- 20.0 Describe principles and perform techniques of preventive dentistry
- 21.0 Perform patient assessment
- 22.0 Perform direct patient services and competently provide dental hygiene process of care for the child, adolescent, adult and geriatric patient as well as the special needs patient
- 23.0 Implement and evaluate community health interventions and research activities

Florida Department of Education
Student Performance Standards

Program Title: Dental Hygiene
CIP Number: 1351060200
Program Length: 88 credit hours
SOC Code(s): 29-2021

The AS degree requires the inclusion of a minimum of 15 credits of general education coursework according to SACS, and it must be transferable according to Rule 6A-14.030 (2), F.A.C. At the completion of this program, the student will be able to:
Dental Hygiene Students completing the following intended outcomes (1- 23) have met the requirements of the Dental Hygiene Program and qualify to make application for the Dental Hygiene National Board and state licensure examinations.

01.0	Demonstrate knowledge of the dental health care delivery system and dental health occupations – The student will be able to:
01.01	Identify the basic components of the dental health care delivery system including public, private, government and non-profit.
01.02	Describe the various types of dental health care providers and the range of services available.
01.03	Describe the composition and functions of a dental health care team
01.04	Identify the general roles and responsibilities of the individual members of the dental health care team.
01.05	Identify the roles and responsibilities of the consumer within the dental healthcare system.
01.06	Explain the cause and effects of factors that influence the current delivery system of dental healthcare.
01.07	Explain the impact of emerging issues including technology, epidemiology, bioethics and socioeconomics on the dental healthcare delivery system.
01.08	Discuss the history of dentistry and dental hygiene
02.0	Use oral and written communication skills in creating, expressing and interpreting information and ideas – The student will be able to:
02.01	Apply basic speaking and active listening skills including reflection, restatement, and clarification techniques.
02.02	Develop basic observational skills and related documentation strategies in written and oral form.
02.03	Identify characteristics of successful and unsuccessful communication including communication styles and barriers.
02.04	Compose written communication using correct spelling, grammar,-formatting and confidentiality and specific formats of letter writing.

02.05	Recognize components of medical and dental terminology and abbreviations.
02.06	Recognize the importance of courtesy and respect for patients and other health care workers and maintain good interpersonal relationships.
02.07	Recognize the importance of patient education regarding dental and health care.
02.08	Adapt communication skills to meet various levels of understanding and orientation of diversity including but not limited to sexual orientation, gender orientation, disability, age, culture, economics, ethnicity and religion.
02.09	Identify psychological considerations influencing communication and behaviors.
03.0	Describe the legal and ethical responsibilities of the dental health care worker – The student will be able to:
03.01	Identify areas of Florida Statute 466 and Rule 64B5-16 FAC and Rule 64B5-25 FAC applicable to practice by the dental health workers.
03.02	Explain practices that could result in malpractice, liability, negligence, abandonment, false imprisonment and fraud.
03.03	Demonstrate procedures for accurate documentation and record keeping.
03.04	Interpret healthcare facility policy and procedures.
03.05	Explain the patients' "Bill of Rights."
03.06	Identify and implement standards of the Health Insurance Portability and Accountability Act (HIPAA).
03.07	Distinguish between express, implied and informed consent.
03.08	Explain the laws governing harassment, labor and employment.
03.09	Differentiate between legal and ethical issues in dentistry.
03.10	Describe a Code of Ethics consistent with the dental hygiene profession.
03.11	Identify and compare personal, professional and organizational ethics.
03.12	Recognize the limits of authority and responsibility of dental health care workers including legislated scope of practice.
03.13	Recognize and report illegal and/or unethical practices of dental health care workers.
03.14	Recognize signs of abuse and neglect.
03.15	Demonstrate an understanding of reporting requirements for all types of abuse including domestic violence and neglect for all ages.
03.16	Identify resources for victims of domestic violence.
03.17	Explain risk management.

04.0	Demonstrate an understanding of general anatomy and physiology and apply wellness and disease concepts – The student will be able to:
04.01	Develop a basic understanding of the structure and function of the body systems.
04.02	Identify common disorders related to each of the body systems.
04.03	Explain basic concepts of positive self-image, wellness, and stress.
04.04	Describe a wellness and stress control plan that can be used in personal and professional life.
05.0	Demonstrate the importance of health, safety, and environmental management systems in dental organizations and their importance to organizational performance and regulatory compliance – The student will be able to:
05.01	Describe personal and jobsite safety rules and regulations that maintain safe and healthy work environments.
05.02	Identify and describe methods in medical error reduction and prevention in the dental healthcare setting.
05.03	Demonstrate an understanding of personal safety procedures based on Occupations Safety and Health Administration (OSHA) and Centers for Disease Control (CDC) regulations (including standard precautions).
05.04	Recognize Safety Data Sheets (SDS) and Globally Harmonized System (GHS) labels and comply with safety signs, symbols and labels.
05.05	Describe procedures for the safe transport and transfer of patients.
05.06	Describe fire safety, disaster and evacuation procedures.
05.07	Explain emergency procedures to follow in response to workplace accidents.
05.08	Demonstrate handwashing and the use of personal protective equipment used in dentistry.
06.0	Recognize and respond to emergency situations – The student will be able to:
06.01	Take and record vital signs.
06.02	Describe legal parameters relating to the administration of emergency care.
06.03	Obtain and maintain training or certification in cardiopulmonary resuscitation (CPR), automated external defibrillator (AED), foreign body airway obstruction (FBAO) and first aid.
07.0	Use information technology tools – The student will be able to:
07.01	Define terms and demonstrate basic computer skills.
07.02	Interpret information from electronic medical documents.
08.0	Explain the importance of employability skills – The student will be able to:
08.01	Identify personal traits or attitudes desirable in a member of the healthcare team.

08.02	Exemplify basic professional standards of dental healthcare workers as they apply to hygiene, dress, language, confidentiality and behavior (i.e. telephone etiquette, courtesy and self-introductions).
08.03	Maintain a career portfolio to document knowledge, skills, and experience.
08.04	Develop a professional resume
08.05	Conduct a job search and complete a job application form correctly.
08.06	Demonstrate effective job interview techniques
08.07	Examine levels of education, credentialing requirements including licensure and certification, employment opportunities, workplace environments and career growth potential.
08.08	Examine licensing, certification, and industry credentialing requirements.
09.0	Demonstrate knowledge of blood borne diseases, including HIV/AIDS – The student will be able to:
09.01	Recognize emerging diseases and disorders.
09.02	Demonstrate knowledge of transmission and treatment of diseases caused by blood borne pathogens including Hepatitis B.
09.03	Identify "at risk" behaviors that promote the spread of diseases caused by blood borne pathogens and the public education necessary to combat the spread of these diseases.
09.04	Identify community resources and services available to the individuals with diseases caused by blood borne pathogens.
09.05	Apply infection control techniques designed to prevent the spread of diseases caused by blood borne pathogens to the care of all patients following Centers for Disease Control (CDC) guidelines.
09.06	Demonstrate knowledge of the legal aspects of treating patients with HIV+ infection and AIDS, including testing.
10.0	Demonstrate leadership and teamwork skills needed to accomplish team goals and objectives – The students will be able to:
10.01	Analyze attributes and attitudes of an effective leader.
10.02	Recognize factors and situations that may lead to conflict.
10.03	Demonstrate effective problem-solving techniques for managing team conflict.
11.0	Perform expanded functions for the dental hygienist as permitted by Florida Statutes/Law--The student will be able to:
11.01	Perform expanded functions as permitted by the Florida Statutes/Law pertaining to the practice of dental hygiene.
12.0	Perform dental office procedures--The student will be able to:
12.01	Maintain appointment control with effective time management skills.
12.02	Assess, create, modify, and maintain an active recare system.

12.03	Prepare and maintain accurate patient records.
12.04	Prepare and maintain dental office inventory control and purchasing.
12.05	Demonstrate skills on office equipment to include computers and dental office management systems.
12.06	Identify correct code on dental procedures and nomenclature (CDT Codes).
12.07	Maintain a positive office environment.
12.08	Receive and dismiss patients and visitors.
12.09	Demonstrate reporting and recording of adverse events.
13.0	Identify, describe, maintain and utilize dental instruments and equipment--The student will be able to:
13.01	Identify various types, functions and operations of dental operatory and laboratory equipment.
13.02	Maintain dental operatory equipment and instruments including proper sharpening techniques.
13.03	Identify types and functions of dental hygiene instruments.
14.0	Identify and perform dental and radiographic procedures--The student will be able to:
14.01	Describe history, physics and biological effects of ionizing radiation.
14.02	Identify parts of the imaging machine including accessories.
14.03	Demonstrate radiologic health protection techniques.
14.04	Perform processing procedures that include application and care.
14.05	Place image receptors and expose dental images that could include chemical emulsion, digital or phosphor plates, and understand the relevance of exposure settings, times and patient record keeping.
14.06	Identify radiographic anatomical landmarks.
14.07	Mount radiographic surveys and/or save and store digital files.
15.0	Identify properties and uses, and manipulate dental materials--The student will be able to:
15.01	Identify properties and uses and manipulation of gypsum.
15.02	Identify properties and uses and manipulation of restorative materials.
15.03	Identify properties and uses and manipulation of dental cements.

15.04	Identify properties and uses and manipulation of impression materials.
15.05	Identify properties and uses and manipulation of acrylics and/or thermoplastics.
15.06	Identify dental laboratory procedures that may include the fabrication of casts, custom trays, temporary crowns and/or bridges.
15.07	Clean removable dental appliances.
16.0	Describe the legal and ethical responsibilities of the dental hygienist--The student will be able to:
16.01	Define commonly used legal vocabulary relating to dentistry.
16.02	Describe ethical considerations/obligations in the dental team-patient relationship.
16.03	Explain risk management and root cause analysis.
16.04	Identify areas of Florida Statute 466 and Rule chapter 64B5 applicable to dentistry and dental hygiene.
16.05	Apply self-assessment skills to prepare for life-long learning.
16.06	Apply ethical principles, legal and regulatory concepts to resolve ethical dilemmas.
17.0	Identify and explain the formation and function of the head, neck, dental structures and tissues including pathological conditions of the human body in relation to the oral cavity--The student will be able to:
17.01	Identify structures and functions of head and neck anatomy including bones, muscles, sinuses, salivary glands, lymph nodes, nerves and blood vessels.
17.02	Identify embryonic development of head, oral cavity, and teeth.
17.03	Describe the histological components of the head, oral cavity, and elements of the teeth and supporting structures.
17.04	Describe and differentiate between normal and malocclusion.
17.05	Identify the elements of the chemical basis of life, cellular metabolism and the structure of the major tissue types of the human body.
17.06	Describe the metabolism of nutrient foods, vitamins and minerals by the human body and pathological conditions related to nutrient deficiencies.
17.07	Identify anatomical structures and physiological function of the principle systems of the human body including the skeletal, muscular, integumentary, circulatory, lymphatic, endocrine, digestive, reproductive, respiratory, urinary, and nervous systems.
17.08	Recognize and describe oral pathological conditions related to the teeth and their supporting structures.
17.09	Identify teeth and their landmarks, and the morphological characteristics of each individual tooth.
17.10	Recognize and describe developmental anomalies related to the teeth, face, and oral structures.
18.0	Identify and explain principles of microbiology, disease transmission, disease prevention, and perform infection control procedures--The

student will be able to:	
18.01	Differentiate between pathogenic and non-pathogenic microorganisms.
18.02	Describe pathogens and modes of disease transmission.
18.03	Differentiate between aseptic and non-aseptic environments.
18.04	Perform aseptic handwashing technique including use of antiseptic gels.
18.05	Describe, apply and differentiate methods of cleaning, disinfection and sterilization
18.06	Recognize the need for and proper precautions for the prevention of disease transmission during all dental related procedures.
18.07	Identify the role of prokaryotic cells, eukaryotic cells, viruses, and bacteria in the infections and mechanisms of diseases.
18.08	Identify the genetics of microbes including replication of DNA and protein synthesis, mutation and gene transfer.
19.0	Identify and explain usage, administration, indications, contraindications, adverse reactions and precautions of pharmaceutical and anesthetic agents used in the treatment of dental disease--The student will be able to:
19.01	Identify drug requirements, agencies, and regulations.
19.02	Record a drug prescription on a patient's chart.
19.03	Identify drug actions, side effects, indications and contraindications; verify with Physician's Desk Reference or its equivalent.
19.04	Describe the process of drug metabolism.
19.05	Identify common drugs used in dentistry.
19.06	Identify pharmaceuticals and medicaments used in the oral cavity.
19.07	Recognize specific conditions in the oral cavity caused by pharmaceutical agents.
19.08	Identify properties of anesthetics.
19.09	Identify the tissues innervated by each of the nerves associated with dental local and topical anesthesia.
19.10	Describe properties and mode of action of an effective local and topical anesthetic.
19.11	List systemic considerations in choosing a local and topical anesthetic.
19.12	Describe methods of administering local and applying topical anesthetics.
19.13	List potential local and systemic adverse reactions associated with local anesthetic administration.

19.14	Prepare armamentarium for administering local anesthetics for recognized techniques.
19.15	Describe the monitoring process and identify precautions in the use of nitrous oxide-oxygen inhalation analgesia.
20.0	Describe principles and perform techniques of preventive dentistry--The student will be able to:
20.01	Identify, communicate, and instruct patients on applicable methods of preventive dentistry that utilize:
20.01.01	risk assessment
20.01.02	evidence based learning
20.01.03	individualized preventive care plans
20.01.04	counseling
20.01.05	training regarding health status and rationale for preventive care plan.
20.02	Identify properties and indications for use of anticariogenic treatments utilized in the community, home, and office.
20.03	Identify and demonstrate proper auxiliary aides based on individual patient needs.
20.04	Identify and describe deficiencies that manifest symptoms in the oral cavity and communicate applicable therapies.
20.05	Formulate and present diets to address specific dental needs and provide nutritional counseling.
21.0	Perform patient assessment--The student will be able to:
21.01	Take, record, and correlate medical/dental history with dental hygiene treatment plan and services to be performed.
21.02	Take, record, and correlate vital sign observations with dental hygiene treatment plan and services to be performed.
21.03	Assess vital signs in order to reduce incidence of patient complications and medical emergencies.
21.04	Perform record and correlate extraoral and intraoral examination findings with dental hygiene treatment plan and patient services to be performed.
21.05	Observe and record existing restorations as well as conditions and suspected pathologies of hard and soft tissues using the appropriate armamentarium.
21.06	Conduct comprehensive periodontal examination including pocket depth, attachment level, recession, mobility, furcations, radiographic findings, and tissue health.
21.07	Consult with dentist and physicians to verify dental and medical information and develop the treatment plan to be implemented.
21.08	Interpret and correlate dental radiographs and dental charting with dental hygiene treatment plan.
21.09	Perform soft tissue reassessment and evaluate the effects of initial dental hygiene therapy and make appropriate therapy modifications or referrals.
21.10	Recognize systemic diseases from oral manifestations.
21.11	Record diagnosis made by dentist.

21.12	Recognize and respond appropriately to contraindications for dental treatment found in medical and dental history.
21.13	Identify and assess dental office emergencies and follow the appropriate protocol for treatment.
22.0	Perform direct patient services and competently provide dental hygiene process of care for the child, adolescent, adult and geriatric patient as well as the special needs patient --The student will be able to:
22.01	Detect calculus for removal and differentiate between deposits and other causes of tooth surface roughness.
22.02	Perform non-surgical periodontal debridement (scaling and root planing) using appropriate armamentarium and instrumentation technique.
22.03	Manipulate mechanical instruments for hard and soft deposit removal, i.e. ultrasonic, air-powder polishing system and/or slow-speed hand-piece.
22.04	Perform oral prophylaxis.
22.05	Demonstrate knowledge of soft tissue curettage.
22.06	Apply desensitizing and/or chemotherapeutic agents where applicable.
22.07	Communicate to patients' appropriate post-operative instructions and correctly select all necessary self-care therapies intended to restore and maintain the individual patient's soft tissue health for long term care.
22.08	Provide and communicate dietary counseling for health maintenance and specific healing needs.
22.09	Provide and communicate recommendations for patient use of caries prevention agents.
22.10	Provide a comprehensive collection of patient data to identify the physical and oral health status as well as risk factors that could affect patient care and healing.
22.11	Provide analysis of assessment findings and use of critical thinking in order to address the patient's dental hygiene treatment needs.
22.12	Establish a dental hygiene care plan that reflects the expected outcomes and treatment interventions to facilitate optimal oral health.
22.13	Present proposed treatment and procedures to the patient and obtain appropriate informed consent signatures prior to rendering patient care services.
22.14	Provide patient-centered treatment and evidence-based care in a manner minimizing risk and optimizing oral health.
22.15	Measure the extent to which expected outcomes identified in the dental hygiene care plan are achieved.
22.16	Complete an accurate recording of all documentation relevant to patient care.
23.0	Implement and evaluate community health interventions and research activities--The student will be able to:
23.01	Demonstrate competence in assessment, planning, implementation and evaluation of community health interventions.
23.02	Formulate and analyze research methodologies for community health interventions.

23.03	Perform a literature search and interpret research findings in scientific literature.
23.04	Apply research findings to dental hygiene care delivery.
23.05	Apply statistical analysis and evidence based research to health trends and community interventions.
23.06	Collaborate and perform a needs assessment with community partners.
23.07	Differentiate scientific value of literature found in both electronic and traditional mediums.

DRAFT

Additional Information

Laboratory Activities

Laboratory investigations that include scientific inquiry, research, measurement, problem solving, emerging technologies, tools and equipment, as well as, experimental, quality, and safety procedures are an integral part of this career and technical program/course. Laboratory investigations benefit all students by developing an understanding of the complexity and ambiguity of empirical work, as well as the skills required to manage, operate, calibrate and troubleshoot equipment/tools used to make observations. Students understand measurement error; and have the skills to aggregate, interpret, and present the resulting data. Equipment and supplies should be provided to enhance hands-on experiences for students.

Equipment and supplies should be provided to enhance hands-on experiences for students. In depth clinical information and requirements can be found in the Commission on Dental Accreditation Dental Hygiene Standards.

Special Notes

General education content must include oral and written communications, Psychology and Sociology.

Biomedical science content must include content in anatomy, physiology, chemistry, biochemistry, microbiology, immunology, general pathology and/or pathophysiology, nutrition and pharmacology.

Dental sciences content must include tooth morphology, head, neck and oral anatomy, oral embryology and histology, oral pathology, radiography, periodontology, pain management, and dental materials.

Graduates must be competent in providing the dental hygiene process of care which includes: Assessment, Planning, Implementation, and Evaluation.

This program meets the Department of Health HIV/AIDS, domestic violence education and prevention of medical errors requirements.

If students in this program are seeking a licensure, certificate or registration through the Department of Health, please refer to 456.0635 F.S. for more information on disqualification for a license, certificate, or registration through the Department of Health.

Outcomes 01-11 are referred to as the Health Careers Core and do not have to be completed if the student has previously completed the Core in another health science program. The Core should be taken first or concurrently with the first course in the program. Following the successful completion of the core, the student is eligible to take the National Health Care Foundation Skill Standards Assessment with instructor approval and the completion of a portfolio.

Career and Technical Student Organization (CTSO)

HOSA: Future Health Professionals is the intercurricular career and technical student organization providing leadership training and reinforcing specific career and technical skills. Career and Technical Student Organizations provide activities for students as an integral part of the instruction offered.

Accommodations

Federal and state legislation requires the provision of accommodations for students with disabilities to meet individual needs and ensure equal access. Postsecondary students with disabilities must self-identify, present documentation, request accommodations if needed, and develop a plan with their counselor and/or instructors. Accommodations received in postsecondary education may differ from those received in secondary education. Accommodations change the way the student is instructed. Students with disabilities may need accommodations in such areas as instructional methods and materials, assignments and assessments, time demands and schedules, learning environment, assistive technology and special communication systems. Documentation of the accommodations requested and provided should be maintained in a confidential file.

Additional Resources

For additional information regarding articulation agreements, Bright Futures Scholarships, Fine Arts/Practical Arts Credit and Equivalent Mathematics and Equally Rigorous Science Courses please refer to:

<http://www.fldoe.org/academics/career-adult-edu/career-tech-edu/program-resources.stml>